

A faunal study of the subfamily Doryctinae in Turkey (Hymenoptera: Braconidae)

Ahmet BEYARSLAN*

Department of Biology, Faculty of Arts and Sciences, Bitlis Eren University, Bitlis, Turkey

Received: 16.02.2014 • Accepted: 08.05.2014 • Published Online: 02.01.2015 • Printed: 30.01.2015

Abstract: From the serial studies of the Braconidae fauna of Turkey, the subfamily Doryctinae was treated. Adult specimens were collected from various habitats of Turkey between 1979 and 2013. In total, 58 species belonging 25 genera were reported for the studied region, among which 52 species were recorded for the first time from Turkey. The numbers of species of each genus are as follows: *Clinocentrus* Haliday, 1833: 1; *Coeloides* Wesmael, 1838: 1; *Colastes* Haliday, 1833: 6; *Dendrosoter* Wesmael, 1838: 1; *Dendrosotinus* Telenga, 1941: 1; *Dolopsidea* Hincks, 1944: 2; *Doryctes* Haliday, 1836: 4; *Ecphyllus* Förster, 1862: 1; *Hecabalodes* Wilkinson, 1929: 1; *Hellenius* Tobias, 1982: 1; *Heterospilus* Haliday, 1836: 7; *Hormius* Nees, 1819: 2; *Leluthia* Cameron, 1887: 4; *Oncophanes* Förster, 1862: 1; *Ontsira* Cameron, 1900: 2; *Pachystigmus* Hellén, 1927: 1; *Pambolus* Haliday, 1836: 1; *Polystenus* Förster, 1862: 1; *Pseudohormius* Tobias & Alexeev, 1973: 1; *Rhaconotus* Ruthe, 1854: 6; *Rhopstrocentrus* Marshall, 1897: 1; *Rhysipolis* Förster, 1862: 4; *Rhyssalus* Haliday, 1833: 1; *Spathiomorpha* Tobias, 1976: 1; and *Spathius* Nees, 1819: 6. The known hosts of each species are given.

Key words: Braconidae, Doryctinae, fauna, new records, Turkey

1. Introduction

The Doryctinae species differ from other cyclostome braconids by the presence of a row of short spines on the anterior edge of the fore tibia, by a flange on the apicolateral edges of the propleuron, and by the apex of the ovipositor with a double node dorsally (Quicke et al., 1992; Marsh, 1993).

Doryctinae is one of the most diversified subfamilies in Braconidae (Shenefelt and Marsh, 1976). More than 1300 doryctine species have been described in 187 recognized genera belonging to 16 tribes, most of which are restricted to tropical and subtropical regions (Belokobylskij, 1992; Yu et al., 2012).

Doryctinae shows a wide range of biological habits, perhaps one of the most diverse in the family Braconidae. Most species for which the biology is known are idiobiont ectoparasitoids of mostly wood-boring beetle larvae in the families Anobiidae, Bostrichidae, Buprestidae, Cerambycidae, Colydiidae, Curculionidae, Eucnemidae, Mordellidae, and Scolitidae (Coleoptera); a few stem-boring lepidopterous larvae in the families Crambidae, Gelechiidae, Gracillariidae, Lyonetiidae, Momphidae, Cosmopterigidae, Bostrichidae, Buccalatricidae, Coleophoridae, Elachistidae, Lymantriidae, Micropterigidae, Nepticulidae, Pyralidae, Tineidae, Tischeriidae, Tortricidae, and Sesiidae (Lepidoptera);

certain sawfly larvae of Cephidae, Tenthredinidae, and Xiphydriidae (Hymenoptera); and Oligochaeta (Lumbricidae) and Aranaea (Theridiidae) (Marsh, 1997; Yu et al., 2012). Although many doryctines attack beetle larvae in seeds, a few species are now known to be phytophagous in seeds or to be gall inducers in various tropical plants (Wharton and Hanson, 2005).

Many taxonomic studies have recently dealt with Doryctinae throughout the world (Fischer, 1970, 1980; Shenefelt and Marsh, 1976; Shaw and Edgerly, 1985; Quicke, 1987, 1997; Belokobylskij, 1990, 1993, 1994a, 1994b, 1995, 1998, 2003, 2004a, 2004b; Ramirez and Marsh, 1996; Marsh, 1997, 2002; Barbalho et al., 1999; Marsh and Melo, 1999; Belokobylskij and Quicke, 2000; van Achterberg, 2003; Belokobylskij and Chen, 2004, 2006; Belokobylskij et al., 2004a, 2004b, 2012; Belokobylskij and Maeto, 2006, 2008, 2009; Gomes and Penteado-Dias, 2007; Belokobylskij and Žikić, 2009; Ghahari, 2009; Kula, 2009; Loni et al., 2012; Martinez and Zaldivar-Riverón, 2013; Tang et al., 2013).

The subfamily Doryctinae is not common in Turkey. The fauna of the Turkish Doryctinae has been poorly studied until recently and only 9 species were recorded so far: *Dendrosoter (Caenopachys) hartigii* (Ratzeburg) by Mancini et al. (2003), *Dendrosoter (D.) middendorffii* (Ratzeburg) by Schimitschek (1941, 1944), *Dendrosoter*

* Correspondence: abeyars@gmail.com

(*D.*) *protuberans* (Nees) by Schimitschek (1939, 1941; Zaldivar-Riverón et al., 2008), *Spathius* (*S.*) *curvicaudis* Ratzeburg by Schimitschek (1944) and Hedqvist (1976), *Spathius* (*S.*) *rubidus* (Rossi) by Belokobylskij (2001), *Pambolus* (*Phaenodus*) *pallipes* (Förster) by Belokobylskij (1986), *Hormius* *propodealis* (Belokobylskij) by Belokobylskij (2001), and *Dolopsidea indagator* (Haliday) and *Dolopsidea tatianae* (Telenga) by Beyarslan and Aydogdu (2013).

D. (*C.*) *hartigii* (Ratzeburg), *D.* (*D.*) *middendorffii* (Ratzeburg), and *H. propodealis* (Belokobylskij) could not be determined in this study.

Adult braconids were collected from various parts of Turkey between 1979 and 2013 and the Doryctinae specimens from this material are studied in this contribution. In total, 58 species belonging to 25 genera are reported for the studied regions of Turkey. Of these, 40 species are distributed in Anatolia (Asian Turkey) and 2 species are distributed only in Thrace (European Turkey), while 16 species occur in both.

2. Materials and methods

Adult specimens of Doryctinae were collected from various habitats of Turkey using Malaise and light traps and sweeping nets. The material is kept in the museum of Bitlis Eren University in Bitlis.

Publications consulted for distribution data include those of Matthews (1970), Belokobylskij and Quicke (2000), van Achterberg (2003), Zaldivar-Riverón and Belokobylskij (2009), and Rastegar et al. (2012).

Relevant literature was used for taxonomical examination of the material. The specimens were identified mostly using the keys of Belokobylskij (1992, 2001, 2003), Belokobylskij and Samartsev (2011), and Tobias (1986). The species of Doryctinae are ectoparasitoids of forest and agricultural pests. Parasitoid doryctine species lay their eggs on the body of the host, which is then used as food for developing larvae. The host is ultimately killed. Hence, a list host species was arranged and is provided in the present study.

Information on parasitoids, hosts, general distributions (in terms of zoogeographical region), and parasitoids of the species determined are given according to Yu et al. (2012). The genera and species are tabulated alphabetically therein.

The terminology and measurements follow van Achterberg (1993). All specimens are deposited in the collection of the Department of Entomology, Bitlis Eren University, Bitlis, Turkey. An asterisk (*) is used within the text for the species that were previously recorded from Turkey.

3. Results

3.1. Genus *Clinocentrus* Haliday, 1833

Clinocentrus vestigator (Haliday, 1836)

Material examined: Erzurum-Olur (39°54'35"N, 41°16'32"E), 1853 m. 12.09.2012, 8♀♀, 3♂♂.

Distribution: Western Palearctic.

Hosts: Lepidoptera, Epermeniidae: *Epermenia chaerophylella* Goeze 1783; *E. illigerella* (Hübner, 1813). Ypsolophidae: *Ypsolopha vittella* (Linnaeus, 1758).

3.2. Genus *Coeloides* Wesmael, 1838

Coeloides forsteri Haeselbarth, 1967

Material examined: Artvin-Merkez (41°36'37"N, 041°35'58"E), 32 m. 06.06.2011 (*Ips sexdentatus*), 1♀.

Distribution: Western Palearctic.

Hosts: Coleoptera, Scolytinae: *Ips typographus* (Linnaeus, 1758). Curculionidae: *Pissodes notatus*; *P. (P.) piceae* (Illiger, 1807); *P. (P.) piniphilus* (Herbst 1797).

3.3. Genus *Colastes* Haliday, 1833

Colastes (*Colastes*) *braconius* Haliday, 1833

Material examined: Bolu-Aladağ-Çepni Yaylası (40°44'0"N, 31°36'0"E), 1470 m. 27.08.2003, 1♀; -Karacasu-Gölcük (40°44'0"N, 31°36'0"E), 1470 m. 26.08.2002, 1♀; Kastamonu-Daday-Ballıdağ (41°28'0"N, 33°25'0"E), 1750 m. 01.07.2001, 5♀♀; Kırklareli-Yeniceköy (41°44'11"N, 27°38'8"E), 500 m. 02.06.1993, 1♀; -Pınarhisar-Sarpdere (41°52'00"N, 27°35'03"E), 192 m. 03.06.1993, 1♀; Trabzon-Mağka-Sümela (40°41'59"N, 39°39'26"E), 1073 m. 03.07.2004, 1♀; 05.08.2005, 1♀; Zonguldak-Devrek-Davulga (41°13'30"N, 31°57'34"E), 800 m. 29.06.2001, 6♀♀.

Distribution: Palearctic. Introduced to USA-Delaware; USA-New Jersey; USA-Pennsylvania (Hendrickson and Barth, 1979).

Hosts: Coleoptera, Curculionidae: *Cryptorhynchus lapathi* (Linnaeus, 1758); *Rynchaenus avellanae* (Donovan, 1797); *R. fagi* (Linnaeus 1758). Diptera, Agromyzidae: *Agromyza albipennis* (Meigen, 1830); *A. frontella* (Rondani, 1875); *A. nana* (Meigen, 1830); *Amauromyza verbasci* (Bouche, 1847); *A. verbasci* (Bouche, 1847); *Aulagromyza cornigera* (Griffiths, 1973); *A. hendeliana* (Hering, 1926); *Chromatomyia aprilina*; (*Goureau*, 1851); *C. loniceriae* (Robineau-Desvoidy, 1851); *L. approximata* (Hendel, 1920); *L. lutea* (Meigen, 1830); *L. pusilla* (Meigen, 1830); *Phytoliriomyza melampyga* (Loew, 1869); *P. variegata* (Meigen, 1830); *Phytomyza aconiti* (Hendel, 1920); *P. affinis* (Fallén, 1823); *P. albiceps* (Meigen, 1830); *P. chaerophylli* (Kaltenbach, 1856); *P. leucanthemi* (Hering, 1935); *P. ranunculi* (Schrank, 1803); *P. xylostei* (Kaltenbach, 1862). Hymenoptera, Tenthredinidae: *Fenus a ulmi* (Sundevall, 1844); *Heterarthrus vagans* (Fallén, 1808); *Messa hortulana* (Klug, 1818); *Pontania bella* (Zaddach, 1876); *P. pedunculi* (Hartig, 1837); *P. proxima* (Serville, 1823). Lepidoptera, Coleophoridae: *Coleophora gryphipennella* (Hübner,

1796). **Cosmopterigidae:** *Cosmopterix scribaiella* (Zeller, 1850). **Elachistidae:** *Elachista humilis* (Zeller, 1850). **Eriocraniidae:** *Dysserocrania subpurpurella* (Haworth, 1828). **Gracillariidae:** *Cameraria ohridella* (Deschka & Dimic, 1986). *Parornix betulae* (Stainton, 1854); *Phyllonorycter acerifoliella* (Zeller, 1839); *P. blancarella* (Fabricius, 1781); *P. cavella* (Zeller, 1846); *P. comparella* (Duponchel, 1843); *P. coryli* (Nicelli, 1851); *P. dubitella* (Herrick-Schäffer, 1855); *P. emberizaepennella* (Bouche, 1834); *P. esperella* (Goeze, 1783); *P. fitchella* (Clemens, 1860); *P. froelichiella* (Zeller, 1839); *P. geniculella* (Ragonot, 1874); *P. harrisella* (Linnaeus, 1761); *P. heegeriella* (Zeller, 1846); *P. insignitella* (Zeller, 1846); *P. lantanella* (Schrank, 1802); *P. lautella* (Zeller, 1846); *P. leucographella* (Zeller, 1850); *P. maestingella* (Müller, 1764); *P. messaniella* (Zeller, 1846); *P. nicellii* (Stainton, 1851); *P. nigrescentella* (Logan, 1851); *P. oxyacantheae* (Frey, 1856); *P. platani* (Staudinger, 1870); *P. platanoidella* (Joannis, 1920); *P. pomonella* (Herrick-Schäffer 1855); *P. populifoliella* (Treitschke, 1833); *P. pyrifoliella* (Gerasimov, 1933); *P. quercifoliella* (Zeller, 1839); *P. quinnata* (Geoffroy, 1785); *P. quinqueguttella* (Stainton, 1851); *P. robiniella* (Clemens, 1859); *P. roboris* (Zeller, 1839); *P. salicolella* (Sircom, 1848); *P. salictella* (Zeller, 1846); *P. saportella* (Zeller, 1839); *P. schreberella* (Fabricius, 1781); *P. sorbi* (Frey, 1855); *P. spinicolella* (Zeller, 1846); *P. stettinensis* (Nicelli, 1852); *P. strigulatella* (Lienig & Zeller, 1846); *P. tenerella* (Joannis, 1915); *P. trifasciella* (Haworth, 1828); *P. Tristrigella* Haworth, 1928; *P. ulmifoliella* (Hubner, 1817); *Antispila treitschkiella* (Fischer von Röslerstamm, 1843); *Heliozela betulae* (Sorhagen, 1885); *Lampides boeticus* (Linnaeus, 1767); *Lyonetia rajella* (Linnaeus 1758); *Mompha epilobiella* (Denis & Schiffermüller, 1775); *Mompha raschiella* (Zeller, 1839). **Nepticulidae:** *Ectoedemia septembrella* (Stainton, 1849); *Stigmella anomalella* (Goeze, 1783); *S. hemargyrella* (Kollar, 1832); *S. microtheriella* (Stainton, 1854); *S. nylandriella* (Tengström, 1848); *S. plagicolella* (Stainton, 1854); *S. ruficapitella* (Haworth, 1828); *Stigmella sorbi* (Stainton, 1861); *Stigmella splendidissimella* (Herrick-Schäffer, 1855); *Stigmella tityrella* (Stainton, 1854). **Pyralidae:** *Pempeliella dilutella* (Denis & Schiffermüller, 1775). **Tischeriidae:** *Emmetia gaunacella* (Duponchel, 1843); *Emmetia heinemanni* (Wocke, 1871); *Emmetia marginea* (Haworth, 1828); *Tischeria dodonaea* (Stainton, 1858); *Tischeria ekebladella* (Bjerkander, 1795); *Selania leplastriana* (Curtis, 1831); *Tortrix viridana* (Linnaeus, 1758); *Ypsolopha dentella* (Fabricius, 1775).

Colastes (Colastes) vividus Papp, 1975

Material examined: Çorum-Dodurga-Osmaniye (40°40'28"N, 34°55'01"E), 1028 m. 28.06.2004, 4♀♀.

Distribution: Western Palearctic.

Hosts: Lepidoptera, Gracillariidae: *Cameraria ohridella* Deschka & Dimic, 1986; *Phyllonorycter spinicolella*

(Zeller, 1846). **Tischeriidae:** *Emmetia heinemanni* (Wocke, 1871).

Colastes (Fungivenator) aciculatus Tobias, 1986

Material examined: Erzincan-Üzümlü-Demirpinar (39°37'38"N, 39°47'34"E), 1200 m. 24.08.2008, 1♂.

Distribution: Western Palearctic.

Colastes (Shawiana) catenator (Haliday, 1836)

Material examined: Ordu-Mesudiye (40°26'45"N, 37°46'42"E), 1330 m. 06.07.2004, 3♀♀.

Distribution: Palearctic.

Hosts: Hymenoptera, Tenthredinidae: *Fenusia dohrni* (Tischbein, 1846); *F. pumila* Leach, 1817; *F. pusilla* (Lepeletier, 1823); *F. rubi* Boie, 1848; *F. ulmi* Sundevall, 1844; *Heterarthrus aceris* (Kaltenbach, 1856); *H. vagans* (Fallén, 1808); *Messa herticulana* (Klug, 1818); *M. nana* (Klug, 1818); *Metallus albipes* (Cameron, 1875); *M. pumilus* (Klug, 1816); *Profenusia pygmaea* (Klug 1816); *Scolioneura betuleti* (Klug, 1816). Lepidoptera, Gracillariidae: *Parna tenella* (Klug, 1816); *Phyllonorycter quercifoliella* (Zeller, 1846). Micropterigidae: *Eriocrania semipurpurella* (Stephens, 1835).

Colastes (Shawiana) laevis (Thomson, 1892)

Material examined: Kars-Sarıkamış-Kayak Tesis-Birinci Etap (40°20'17"N, 42°34'23"E), 2102 m. 19.09.2012, 3♀♀.

Distribution: Palearctic and Oriental.

Hosts: Hymenoptera, Tenthredinidae: *Euura atra* (Jurine 1807); *Euura mucronata* Hartig, 1837; *Fenusia dohrnii* (Tischbein); *Heterarthrus aceris* (Kaltenbach, 1856); *H. microcephalus* (Klug, 1818); *H. vagans* (Fallén, 1808); *Phyllotoma wuestneii* Konow, 1905; *Pontania bella* (Zaddach 1876); *Pontania vesicator* (Bremi-Wolf, 1849); *Tenthredo betuleti* Klug, 1816; *Scolioneura vicina* (Konow, 1894).

Colastes (Xenarcha) lustrator (Haliday, 1836)

Material examined: Aksaray-Ihlara-Yaprakhisar (38°13'18"N, 34°17'14"E), 1354 m. 11.09.2006, 1♀; Ardahan-Posof (41°30'30"N, 42°44'03"E), 1445 m. 21.08.2013, 2♀♀; Amasya-Taşova-Keçiyağı (40°48'49"N, 36°09'41"E), 1066 m. 08.07.2003, 1♀; Bilecik-Küpü (40°05'50"N, 30°00'16"E), 500 m. 20.09.1992, 1♀; Bilecik-Başköy (40°5'32"N, 30°0'22"E), 62 m. 20.09.1992, 1♀; Bolu-Aladağ-Kökez (40°43'50"N, 31°36'34"E), 1000 m. 27.08.2002, 1♀; Bursa-İnegöl (40°4'59"N, 29°30'59"E), 400 m. 06.06.2011, 1♀; Erzurum-İspir-Çamlıkaya (40°37'49"N, 41°10'52"E), 950 m. 16.08.2013, 1♀; -İspir-Yeşilyurt köyü (40°31'29"N, 41°04'55"E), 1570 m. 26.06.2013, 1♀, 1♂; -Olur (39°54'35"N, 41°16'32"E), 1853 m. 12.09.2012, 1♀; -Şenkaya-Gözalan (40°44'36"N, 42°17'40"E), 1500 m. 13.09.2012, 2♀♀; Kastamonu-Ilgaz Dağı (41°06'49"N, 33°55'08"E), 920 m. 30.08.2002, 2♀♀; -Akkaya (41°13'37"N, 33°28'7"E), 110 m. 31.08.2002,

1♀; Bolu-Aladağ-Kökez ($40^{\circ}43'50''N$, $31^{\circ}36'34''E$), 1000 m. 27.08.2002, 1♀; Samsun-Kavrak-Biçi ($41^{\circ}7'24''N$, $36^{\circ}8'32''E$), 19 m. 02.07.2003, 1♀; -Salıpazarı ($41^{\circ}49'98''N$, $36^{\circ}49'35''E$), 650 m. 03.07.2003, 1♀.

Distribution: Western Palearctic.

Hosts: Hymenoptera, Tenthredinidae: *Fenella nigrita* Westwood 1839; *Fenusia pumila* Leach, 1817; *F. pusilla* (Lepeletier, 1823); *F. (Kaliofenusia) ulmi* Sundevall 1844; *Metallus albipes* (Cameron, 1875); *M. pumilus* (Klug, 1816). Lepidoptera, Coleophoridae: *Coleophora*

3.4. Genus *Dendrosoter* Wesmael, 1838

**Dendrosoter (Dendrosoter) protuberans* (Nees, 1834)

Material examined: Edirne Trakya Üniv. Kampüs ($41^{\circ}38'20''N$, $26^{\circ}36'53''E$), 41 m. 24.07.1989, 1♀; Erzincan ($40^{\circ}01'28''N$, $40^{\circ}10'23''E$), 1185 m. 26.08.2008, 1♀; İstanbul-Sarıyer-Bilezikçi çiftliği ($41^{\circ}10'53''N$, $29^{\circ}2'18''E$), 35 m. 14.06.2001, 1♀; Sivas-Koyulhisar ($40^{\circ}17'11''N$, $37^{\circ}49'05''E$), 694 m. 01.07.2004, 1♀; Sivas-Ulaş-Yağdonduran ($39^{\circ}20'28''N$, $37^{\circ}08'43''E$), 1630 m. 01.06.2007, 1♀.

Distribution: Holarctic.

Hosts: Coleoptera, Buprestidae: *Agrilus sulcicollis* Lacordaire, 1835; *Anthaxia hackeri* Frivaldszki, 1884. Cerambycidae: *Exocentrus lusitanus* (Linnaeus, 1767); *Phymatodes testaceus* (Linnaeus, 1758); *Pyrrhidium sanguineum* (Linné, 1758); *Stromatium fulvum* Linsley & Chemsak, 1997; *S. unicolor* (Olivier, 1795). Chrysomelidae: *Agelastica alni* (Linnaeus, 1758). Curculionidae: *Dendroctonus minor* Hartig 1834; *Blastophagus piniperda* (Linnaeus, 1758); *Ceutorhynchus quadridens* Panzer, 1795; *Chaetoptelius vestitus* (Mulsant & Rey, 1860); *C. vestitus* Fuchs, 1913; *Dryocoetes villosus* (Fabricius 1792); *D. villosus* (Fabricius, 1792); *Hylesinus crenatus* (Fabricius, 1787); *Bostrichus fraxini* Panzer, 1779; *B. suturalis* Gyllenhal, 1827; *Hylesinus fraxini* (Panzer, 1779); *H. oleiperda* Fabricius, 1801; *Ips sexdentatus* (Börner, 1767); *I. typographus* (Linnaeus, 1758); *Leperisinus varius* (Fabricius, 1775); *Phloeosinus armatus* Reitter, 1887; *P. aubei* (Perris, 1855); *P. bicolor* (Brulle, 1832); *P. thujae* (Perris, 1855); *Phloeotribus scarabaeoides* (Bernard, 1788); *Pityogenes bidentatus* (Herbst, 1784); *P. quadridens* (Härtig, 1834); *Pteleobius kraatzi* (Eichhoff, 1864); *Scolytus aceris* Knotek, 1892b; *Scolytus amygdali* Guerin, 1847; *S. carpini* (Ratzeburg, 1837); *S. intricatus* (Ratzeburg, 1837); *S. kirschi* Skalitzky, 1876; *S. koenigi* Schevyrew 1890; *S. multistriatus* (Marsham, 1802); *S. pygmaeus* (Fabricius, 1787); *S. ratzeburgi* Janson, 1856; *S. rugulosus* (Mueller, 1818); *S. scotylus* (Fabricius 1775); *Xyleborus dispar* (Fabricius, 1792); *X. dryographus* (Ratzeburg, 1837).

3.5. Genus *Dendrosotinus* Telenga, 1941

**Dendrosotinus ferrugineus* (Marshall, 1888)

Material examined: Ardahan-Altaş ($41^{\circ}9'50''N$, $42^{\circ}52'45''E$), 1800 m. 14.09.2012, 1♀; Balıkesir-Kuş Cenneti ($40^{\circ}2'47''N$, $27^{\circ}58'11''E$), 15 m. 11.05.1991, 1♀; Bartın-Amasra-Kalaycı ($41^{\circ}47'03''N$, $32^{\circ}31'47''E$), 15 m. 27.05.2007, 2♀♀; Erzincan-Tercan-Mercan ($39^{\circ}46'08''N$, $40^{\circ}18'21''E$), 1398 m. 24.08.2008, 1♀; Erzurum-İspir-Yeşilyurt köyü ($40^{\circ}31'29''N$, $41^{\circ}04'55''E$), 1570 m. 26.06.2013, 1♀; İğdır-Karakoyun-Gökçeli ($39^{\circ}59'09''N$, $44^{\circ}10'16''E$), 850 m. 21.06.2013, 1♀; İğdır-Tuzluca-Gaziler ($40^{\circ}06'36''N$, $43^{\circ}25'31''E$), 1035 m. 27.06.2012, 1♀; Kars-Sarıkamış-Şht. Kr. Alb. F. Sungur Tplm. Güzergahı ($40^{\circ}21'44''N$, $42^{\circ}32'32''E$), 2530 m. 25.06.2013, 1♂; Kayseri-Erciyes ($38^{\circ}36'56''N$, $35^{\circ}30'59''E$), 1701 m. 25.06.2009, 1♀; Konya-Çumra-Avdul ($37^{\circ}25'26''N$, $32^{\circ}44'51''E$), 1073 m. 27.06.2009, 1♀; Muğla-Milas-Sarıçay ($37^{\circ}18'36''N$, $27^{\circ}46'48''E$), 50 m. 30.07.1997, 1♀.

Distribution: Western Palearctic.

Hosts: Coleoptera, Cerambycidae: *Scobicia chevrieri* (Villa & Villa, 1835); *Sinoxylon sexdentatum* (Olivier, 1790). Curculionidae: *Phloeotribus scarabaeoides* (Bernard 1788). Scolytidae: *Chaetoptelius vestitus* (Mulsant & Rey 1860).

3.6. Genus *Dolopsidea* Hincks, 1944

**Dolopsidea indagator* (Haliday, 1836)

Material examined: Erzurum-İspir-Çamlıkaya ($40^{\circ}37'49''N$, $41^{\circ}10'52''E$), 950 m. 16.08.2013, 1♂; -İlica ($39^{\circ}56'23''N$, $41^{\circ}06'05''E$), 1895 m. 19.06.2012, 1♀; -Şenkaya-İkizpinar ($40^{\circ}36'28''N$, $42^{\circ}21'00''E$), 1700 m. 27.06.2013, 1♂; -Tortum-Serdarlı ($40^{\circ}28'0''N$, $41^{\circ}18'0''E$), 1685 m. 19.06.2012, 1♀; İğdır-Tuzluca ($40^{\circ}02'51''N$, $43^{\circ}42'54''E$), 1038 m. 23.08.2013, 1♂; Sivas-Zara-Bulakbaşı ($39^{\circ}52'46''N$, $37^{\circ}33'31''E$), 1297 m. 21.05.2007, 1♀.

Distribution: Palearctic.

Hosts: Coleoptera, Anobiidae: *Dorcatoma chrysomelina* Sturm, 1837; *D. dresdensis* Herbst, 1792; Cerambicyidae: *Pogonocherus hispidus* (Linné, 1758).

**Dolopsidea tatianae* (Telenga, 1941)

Material examined: Sivas-Gürün ($38^{\circ}46'05''N$, $37^{\circ}06'03''E$), 1621 m. 05.06.2002, 2♀♀.

Distribution: Western Palearctic.

3.7. Genus *Doryctes* Haliday, 1836

**Doryctes (Doryctes) leucogaster* (Nees, 1834)

Material examined: Adana-Yumurtalık ($36^{\circ}45'59''N$, $35^{\circ}46'59''E$), 13 m. 19.03.2007, 1♀; Amasya-Gümüş-Hacıköy ($40^{\circ}53'25''N$, $35^{\circ}12'07''E$), 849 m. 07.04.2004, 1♀; Artvin-Varziya ($41^{\circ}10'54''N$, $41^{\circ}49'11''E$), 345 m. 03.09.2004, 1♀; -Yusufeli-Küplice ($40^{\circ}56'06''N$, $40^{\circ}30'23''E$), 828 m. 04.09.2004, 1♀; Aydın-Germencik ($37^{\circ}52'19''N$, $27^{\circ}36'6''E$), 70 m. 04.07.2000, 1♀;

Çanakkale-Anzak Mezarlığı ($40^{\circ}15'41''N$, $26^{\circ}16'51''E$), 73 m. 06.05.1993, 1♀; -Ayvacık ($39^{\circ}36'4''N$, $26^{\circ}24'16''E$), 350 m. 23.07.1989, 1♀; Çorum-Kargı ($41^{\circ}13'44''N$, $34^{\circ}48'72''E$), 110 m. 31.08.2002, 1♀; Denizli-Buharkent (*H. griseus*) ($37^{\circ}57'42''N$, $28^{\circ}44'32''E$), 173 m. 11.09.2000, 2♀; -Kale-Alanyurt ($37^{\circ}20'29''N$, $28^{\circ}48'30''E$), 490 m. 08.04.1998, 1♂; Edirne-Lalapaşa-Hacidanışment ($41^{\circ}54'35''N$, $26^{\circ}49'17''E$), 210 m. 05.07.1997, 1♀; Elazığ-Keban-Aşgören ($38^{\circ}47'38''N$, $38^{\circ}44'30''E$), 780 m. 04.06.2007, 1♀; Erzincan-Kemaliye-Yuva ($39^{\circ}15'12''N$, $38^{\circ}30'10''E$), 830 m. 26.08.2008, 1♀; Erzurum-Aşkale ($39^{\circ}55'17''N$, $40^{\circ}42'24''E$), 1670 m. 18.8.2013, 1♀; Eskişehir-A.Ü. Hatıra Ormanı ($39^{\circ}43'09''N$, $30^{\circ}40'37''E$), 907 m. 10.07.2007, 1♀; -Sivrihisar-Babatat ($39^{\circ}31'42''N$, $31^{\circ}37'36''E$), 868 m. 10.07.2007, 1♀, 3♂♂; -Bilecik yolu 26 km ($39^{\circ}55'50''N$, $30^{\circ}16'32''E$), 839 m. 07.07.2007, 2♀♀, 11♂♂; Isparta-Yalvaç ($38^{\circ}17'44''N$, $31^{\circ}10'48''E$), 1150 m. 10.07.2002, 1♀; Kars-Kağızman-Aydınkavak ($40^{\circ}09'11''N$, $43^{\circ}19'14''E$), 1115 m. 20.09.2012, 1♀; Kastamonu-Araç-Akgeçit ($41^{\circ}13'13''N$, $33^{\circ}12'25''E$), 400 m. 28.06.2002, 1♀; Kırklareli-Demirköy-Balaban ($41^{\circ}49'40''N$, $27^{\circ}40'44''E$), 390 m. 15.9.1999, 1♂; -Dereköy ($41^{\circ}56'4''N$, $27^{\circ}21'57''E$), 500 m. 18.06.1986, 1♀; Rize-Ayder Yaylası ($40^{\circ}57'12''N$, $41^{\circ}06'05''E$), 1260 m. 02.09.2004, 1♀; Sivas-Koyulhisar ($40^{\circ}17'11''N$, $37^{\circ}49'05''E$), 694 m. 01.07.2004, 1♀; -Suşehri-Yaygınsoğut ($40^{\circ}16'10''N$, $38^{\circ}09'15''E$), 834 m. 20.08.2004, 1♀; Şanlıurfa-Bozova-Yaslıca mey. ($37^{\circ}26'21''N$, $38^{\circ}23'10''E$), 519 m. 16.06.2007, 1♀, 1♀; 30.04.2006, 2♀♀; -Hilvan-Faik ($37^{\circ}35'19''N$, $38^{\circ}57'19''E$), 585 m. 14.06.2007, 3♀♀; Tekirdağ-Işıklar-Bakıncık ($40^{\circ}48'54''N$, $27^{\circ}22'50''E$), 300 m. 25.08.2000, 1♀; -Uçmakdere ($40^{\circ}47'51''N$, $27^{\circ}21'56''E$), 110 m. 24.08.2000, 1♀; Yozgat-Kuşçukköy ($39^{\circ}39'15''N$, $34^{\circ}43'14''E$), 1202 m. 29.05.2007, 3♂♂.

Distribution: Palearctic, Oriental.

Hosts: Coleoptera, Anobiidae: *Coelostethus pertinax* LeConte, 1861; *Anobium punctatum* (De Geer, 1774); *Anthaxia* sp. Bostrichidae: *Bostrichus capucinus* (Linnaeus, 1758). Buprestidae: *Agrilus* (*Anambus*) *biguttatus* (Fabricius, 1777); *Buprestis aurulenta* Linnaeus, 1767 (syn. *Anthaxia aurulenta*); *Anthaxia* (*Anthaxia*) *manca* (Linnaeus, 1767); *Lampra mirifica* Kraatz, 1880. Cerambycidae: *Acanthocinus aedilis* (Linnaeus, 1758); *Acanthoderes cinereus* Fabricius; *Chlorophorus pilosus* (Forster, 1771); *Chrysobothris salieri* Castelnau & Gory, 1893; *Exocentrus lusitanus* Linnaeus 1767; *Hesperophanes cinereus* Aurivillius, 1912; *Hylotrupes bajulus* (Linnaeus, 1758); *Penichroa fasciata* (Stephens, 1831); *Phymatodes testaceus* (Linnaeus, 1758); *Plagionotus arcuatus* (Linnaeus, 1758); *Rhagium (Hagrium) bifasciatum* Fabricius, 1775; *Rhagium (Rhagium) inquisitor* (Linnaeus, 1758); *Rhagium mordax* (De Geer, 1775); *Tetropium castaneum* (Linnaeus, 1758). Chrysomelidae: *Agelastica alni* (Linnaeus, 1758).

Crambidae: *Loxostege sticticalis* (Linnaeus, 1761). Lepidoptera, Lymantriidae: *Lymantria dispar* Linnaeus, 1758. Aranidae: Theridiidae: *Opilo domesticus* (Sturm, 1837).

Doryctes (Doryctes) molorchi Fischer, 1971

Material examined: Eskişehir-Sündiken Dağı-Karakütük ($39^{\circ}58'00''N$, $31^{\circ}06'00''E$), 788 m. 05.09.2006, 1♀.

Distribution: Western Palearctic.

Hosts: Coleoptera, Cerambycidae: *Molorchus umbellatarum* (Schreber, 1759).

Doryctes (Doryctes) striatellus (Nees, 1834)

Material examined: Elazığ- Keban-Piran ($38^{\circ}47'38''N$, $38^{\circ}44'30''E$), 780 m. 11.06.2008, 5♀♀.

Distribution: Palearctic.

Hosts: Coleoptera, Anobiidae: *Anobium abietis* (Fabricius, 1792); *Dorcatoma dresdensis* Herbst, 1792; *Ernobius mollis* (Linnaeus, 1758). Buprestidae: *Anthaxia tuerki* Ganglbauer, 1886; *Agrilus biguttatus* (Fabricius, 1777). Cerambycidae: *Callidium violaceum* (Linné, 1758); *C. abdominalis* (Olivier, 1795); *Obrium brunneum* (Fabricius, 1792); *Pogonocherus hispidus* (Linnaeus, 1758); *Pyrrhidium sanguineum* (Linnaeus, 1758); *Saperda similis* (Laicharting, 1784); *Tetropium castaneum* (Linné, 1758). Curculionidae: *Hylurgus ligniperda* (Fabricius, 1787); *Magdalais (Magdalais) violacea* (Linnaeus, 1758); *M. rufa* Germar, 1824; *Pissodes notatus* Fabricius, 1787. Scolytidae: *Hylurgops palliatus* (Gyllenhal, 1813); *Ips typographus* (Linnaeus, 1758); *S. ratzeburgi* Janson, 1856; *Tetropium fuscum* (Fabricius, 1787); *T. gabrieli* Weise, 1905. Hymenoptera, Xiphydriidae: *Xiphydria camelus* (Linne, 1758); *Xiphydria prolongata* (Geoffroy, 1785). Lepidoptera, Tortricidae: *Cydia pomonella* (Linnaeus, 1758).

Doryctes (Doryctes) undulatus (Ratzeburg, 1852)

Material examined: Gümüşhane-Şiran-Yeşilbüük ($40^{\circ}13'55''N$, $38^{\circ}58'47''E$), 1434 m. 29.08.2004, 1♀.

Distribution: Palearctic.

Host: Coleoptera, Buprestidae: *Agrilus convexicollis* Redtenbacher, 1849; *A. cuprescens* (Ménétriés, 1832); *A. mendax* (Mannerheim, 1837); *A. viridis* (Linnaeus, 1758); *Anthaxia tuerki* Ganglbauer 1886; *Axinopalpis gracilis* (Krynicki, 1832); *Grammoptera ruficornis* (Fabricius, 1781); *Magdalais armigera* (Fourcroy, 1785); *M. ruficornis* (Linnaeus, 1758). Cerambycidae: *Molorchus umbellatarum* (Schreber, 1759); *M. kiesenwetteri* (Mulsant & Rey, 1861); *Pogonocherus decoratus* Fairmaire, 1855; *P. fasciculatus* (De Geer, 1775); *P. hispidulus* (Piller & Mitterpacher, 1783); *P. hispidus* (Linnaeus, 1758); *Tetrops praeustus* (Linnaeus, 1758). Scolytinae: *Pityogenes bidentatus* (Herbst, 1783).

3.8. Genus *Ecphylus* Förster, 1862

Ecphylus (Ecphylus) silesiacus (Ratzeburg, 1848)

Material examined: Balikesir-Havran (39°33'30"N, 27°5'53"E), 44 m. 20.09.2004, 1♀; Elazığ-Baskil-Canbeyler (38°33'24"N, 38°49'00"E), 1116 m. 03.06.2007, 1♂; -Baskil-Hacimustafa (38°38'34"N, 38°51'07"E), 1436 m. 16.07.2007, 1♂; Tekirdağ-Ganos (40°47'00"N, 27°16'00"E), 550–640 m. 14.07.1986, 1♂; Ankara (40°1'15"N, 32°49'51"E), 870 m. 08.08.2010, 1♀; Malatya-Tekev (38°21'18"N, 38°18'34"E), 977 m. 04.08.1997, 1♀.

Distribution: Palearctic.

Hosts: Coleoptera; Bostrichidae: *Bostrichus laricis* Fabricius, 1792. Curculionidae: *Blastophagus minor* (Hartig, 1834); *Bostrichus binodulus* Ratzeburg, 1837; *Carpoborus minimus* (Fabricius 1798); *Ceutorhynchus pallidactylus* (Marsham 1802); *C. quadridens* Panzer, 1795; *Chaetoptelius vestitus* (Mulsant & Rey 1860); *Cryphalus abietis* (Ratzeburg, 1837); *Ernroporus tilliae* (Panzer, 1793); *Tomicus minor* (Hartig, 1834); *Trypophloeus asperatus* (Gyllenhal, 1813); *Cryphalus binodulus* (Ratzeburg, 1837); *C. piceae* (Ratzeburg, 1837); *C. cinereus* (Herbst, 1793); *Dryocoetes alni* (Georg, 1856); *D. villosus* (Fabricius, 1792); *Ernporicus fagi* (Fabricius, 1798); *Hylesinus crenatus* (Fabricius, 1787); *H. fraxini* (Panzer 1779); *H. spartii* Noerdlinger, 1847; *Ips acuminatus* (Gyllenhal, 1827); *I. typographus* (Linnaeus, 1758); *Liparthrum colchicum* Semenov, 1903; *Magdalais (Odontomagdalais) armigera* (Geoffroy, 1785); *M. (Porrothus) cerasi* (Linnaeus, 1758); *M. (Edo) ruficornis* (Linnaeus, 1758); *Miarus graminis* (Gyllenhal, 1813); *Orthotomicus laricis* (Fabricius, 1792); *Phloeotribus caucasicus* Reitter, 1891; *P. hercegovinensis* (Seitner, 1920); *P. rhododactylus rhododactylus* (Marsham, 1802); *P. scarabaeoides* (Bernard, 1788); *P. spinulosus* (Rey, 1883). Scolytinae: *Pityogenes bidentatus* (Herbst, 1784); *P. bistridentatus* (Eichhoff, 1878); *P. calcaratus* (Eichhoff, 1878); *P. chalcographus* (Linnaeus, 1761); *P. quadridens* (Hartig, 1834); *P. spinidens* (Reitter, 1894); *P. vorontzovi* (Jacobson, 1895); *Pityophthorus micrographus* (Linnaeus, 1758); *P. pityographus* (Ratzeburg, 1837); *Polygraphus poligraphus* (Linnaeus, 1758); *P. subopacus* Thomson, 1871; *Pteleobius kraatzi* (Eichhoff, 1864); *P. vittatus* (Fabricius, 1787); *Scolytus* sp.; *S. carpini* (Ratzeburg, 1837); *S. ensifer* Eichhoff, 1881; *S. intricatus* (Ratz., 1837); *S. kirschi* Skalitzky, 1876; *S. laevis* Chapuis, 1869; *S. mali* (Bechstein, 1805); *S. multistriatus* (Marsham, 1802); *S. pygmaeus* (Fabricius, 1787); *S. rugulosus* (Müller, 1818); *S. scolytus* (Fabricius, 1775). Curculionidae: *Trypodendron domesticum* (Linnaeus, 1758); *Trypophloeus asperatus* (Gyllenhal, 1813); *Xyleborinus saxeseni* (Ratzeburg, 1837); *X. pilosus* (Scopoli, 1763).

3.9. Genus *Hecabalodes* Wilkinson, 1929

Hecabalodes radialis Tobias, 1962

Material examined: Ankara-Temelli-Alagöz (39°45'18"N, 32°26'42"E), 803 m. 10.07.2007, 2♀♀; Aydin-Adnan Menderes Üniv.-Ziraat Fak. (37°50'39"N, 27°50'44"E), 70 m. 18.09.1996, 1♀; Edirne-Trakya Üniv.-Güllapoğlu (41°38'20"N, 26°36'53"E), 41 m. 03.10.2001, I.Tuzağı, 1♀; Elazığ-Keban-İbolar (38°49'03"N, 38°52'16"E), 888 m. 10.07.2007, 1♀; -Keban-Ulupınar (38°44'59"N, 38°51'57"E), 1229 m. 11.06.2008, 1♂; -Merkez (38°40'15"N, 39°11'16"E), 1067 m. 30.05.2003, 4♀♀; Erzincan-İliç-Bağışdaş (39°26'46"N, 38°29'38"E), 1090 m. 26.08.2008, 1♀; -Kemah-Soğukpinar (39°36'06"N, 39°02'17"E), 1040 m. 26.08.2008, 1♀; Eskişehir-A.Ü-Hatıra Ormanı (39°43'09"N, 30°40'37"E), 907 m. 10.07.2007, 1♂; İğdır-Tigem (39°47'10"N, 44°37'05"E), 850 m. 22.06.2013, 1♂; Karabük-Bostanbüyük (41°12'53"N, 32°41'21"E), 290 m. 13.06.2002, 1♀; Malatya-Sultansuyu Harası (38°20'36"N, 38°03'35"E), 881 m. 14.07.2007, 1♀; Mardin-Merkez (37°18'44"N, 40°44'6"E), 1083 m. (Buprestidae host), 10.05.2004, 2♀♀, 1♂; Muğla-Yatağan-Dipsiz (37°25'00"N, 28°06'00"E), 376 m. 31.07.1997, 1♀; Nevşehir-Avanos-Saruhan (38°42'47"N, 34°54'47"E), 958 m. 06.06.2007, 1♀; Samsun-Havza-Mismiliağaç (40°59'51"N, 35°41'49"E), 610 m. 02.07.2003, 1♀; Vezirköprü-Erduası (41°21'27"N, 35°03'06"E), 110 m. 02.07.2003, 1♀; Sinop-Ayancık-B.Zaviye (41°56'49"N, 34°35'30"E), 16 m. 06.09.2001, 1♀; -Yeniçam (41°96'5"N, 34°85'6"E), 30 m. 02.07.2001, 1♀. Tekirdağ-Murathî (41°10'26"N, 27°30'31"E), 60 m. 26.08.1992, 1♀; Çanakkale-Tigem (40°9'19"N, 26°24'50"E), 10 m. 06.07.1996, 1♀; Yozgat-Şefaath-Karanlıkdere (39°34'28"N, 34°38'48"E), 840 m. 29.05.2007, 1♀.

Distribution: Palearctic, Oriental.

Hosts: Coleoptera, Buprestidae: *Acmaeoderidae* sp.; *Acmaeoderella glasunovi* (Semenov, 1895); *Anthaxia* sp.; *Anthaxia judinae* Stepanov, 1954; *Magdalais egregia* Faust, 1885; *M. myochroa* Reichardt, 1928; *Scolytus gretschkini* Sokanovskii, 1956; *Turanium scabrum* (Kraatz, 1882).

3.10. Genus *Hellenius* Tobias, 1982

Hellenius (Hellenius) semiruber (Hellén, 1958)

Material examined: Sivas-Gürün-İncesu (38°46'05"N, 37°06'03"E), 1621 m. 23.06.2009, 1♀.

Distribution: Western Palearctic.

3.11. Genus *Heterospilus* Haliday, 1836

Heterospilus (Heterospilus) cephi Rohwer, 1925

Material examined: Ankara-Temelli-Alagöz (39°45'18"N, 32°26'42"E), 803 m. 17.07.2007, 1♀; Aydin-Söke-Sarıkemer (37°33'42"N, 27°22'8"E), 38 m. 31.07.1997, 1♀; Edirne-Budak-Doğanca (41°45'40"N, 26°20'26"E), 45 m. 30.05.1993, 1♀; -Lalapaşa (41°50'24"N, 26°43'48"E), 560 m. 24.05.1992, 1♀; -Suakacağı (41°50'25"N, 26°35'18"E), 210 m. 05.07.1997, 1♀; Elazığ-Aydınlar

($38^{\circ}48'04''N$, $38^{\circ}53'40''E$), 985 m. 10.07.2007, 1♀; -Baskılı-Canbeyler ($38^{\circ}33'24''N$, $38^{\circ}49'00''E$), 1116 m. 03.06.2007, 1♀; -Keban-İbolar ($38^{\circ}49'03''N$, $38^{\circ}52'16''E$), 888 m. 10.07.2007, 1♀; Eskişehir-Mihallicik-Çatacık ($39^{\circ}58'07''N$, $31^{\circ}10'38''E$), 1332 m. 06.09.2006, 1♀; Gümüşhane-Kelkit-Köycük ($40^{\circ}9'1''N$, $39^{\circ}25'19''E$), 1450 m. 08.08.2005, 1♀; İstanbul-Bahçeköy-Bilezikçi Çiftliği ($41^{\circ}11'33''N$, $28^{\circ}59'53''E$), 30 m. 24.06.1993, 1♀; Malatya-Akçadağ-Darıca ($38^{\circ}20'39''N$, $37^{\circ}40'22''E$), 1540 m. 05.06.2007, 1♀; -Akçadağ-Kürecik ($38^{\circ}20'49''N$, $37^{\circ}46'10''E$), 1749 m. 07.07.2008, 1♀; -Hekimhan ($38^{\circ}49'44''N$, $37^{\circ}55'00''E$), 1094 m. 01.06.2007, 1♀; -Sultansuyu Harası ($38^{\circ}20'36''N$, $38^{\circ}03'35''E$), 881 m. 14.07.2007, 1♀; Muğla-Ula-Yaylasöğüt ($37^{\circ}9'43''N$, $28^{\circ}33'49''E$), 700 m. 30.07.1997, 1♀; -Yatağan ($37^{\circ}20'26''N$, $28^{\circ}8'25''E$), 419 m. 31.07.1997, 1♀; Tekirdağ-Işıklar-Bakıncak ($40^{\circ}48'54''N$, $27^{\circ}22'50''E$), 300 m. 25.08.2000, 1♀.

Distribution: Holarctic.

Hosts: Hymenoptera, Cephidae: *Cephus pygmeus* Latreille, 1803; *Trachelus tabidus* (Fabricius, 1775).

Heterospilus (Heterospilus) genalis Tobias, 1976

Material examined: Malatya-Sultansuyu Harası ($38^{\circ}20'36''N$, $38^{\circ}03'35''E$), 881 m. 14.07.2007, 1♀.

Distribution: Western Palearctic.

Heterospilus (Heterospilus) hemipterus (Thomson, 1892)

Material examined: Tekirdağ-Ganos Dağı ($40^{\circ}47'00''N$, $27^{\circ}16'00''E$), 640 m. 14.07.1986, 1♀.

Distribution: Palearctic.

Heterospilus (Heterospilus) leptosoma Fischer, 1960

Material examined: Adapazarı-Hendek-Hüseyinşeyh ($37^{\circ}20'26''N$, $28^{\circ}8'25''E$), 240 m. 27.06.2001, 1♀; Antalya-Merkez-Patlıcan ($36^{\circ}30'46''N$, $32^{\circ}5'9''E$), 48 m. 09.11.1995, 1♀; İstanbul-Sarıyer-Bahçeköy ($41^{\circ}11'33''N$, $28^{\circ}59'53''E$), 120 m. 14.07.2002, 1♀; Karabük-Bostanbüyük ($41^{\circ}12'53''N$, $32^{\circ}41'21''E$), 290 m. 28.08.2002, 2♀♀; -E.Pazar-Ortaköy ($40^{\circ}55'6''N$, $32^{\circ}34'30''E$), 870 m. 28.08.2002, 1♀.

Distribution: Palearctic, Oriental.

Heterospilus (Heterospilus) rubicola Fischer, 1968

Material examined: Artvin-Yusufeli-Küplüce ($40^{\circ}56'06''N$, $40^{\circ}30'23''E$), 828 m. 01.07.2005, 1♀; -Yusufeli-Küplüce ($40^{\circ}56'06''N$, $40^{\circ}30'23''E$), 828 m. 01.07.2005, 1♀; Bartın-Merkez ($41^{\circ}38'4''N$, $32^{\circ}20'15''E$), 10 m. 09.06.2002, 1♀; -Merkez ($41^{\circ}38'4''N$, $32^{\circ}20'15''E$), 10 m. 09.06.2002, 1♀; Bilecik-Çakırpınar ($40^{\circ}25'11''N$, $29^{\circ}15'20''E$), 795 m. 09.07.1993, 1♀; Gümüşhane-Kelkit-Yeniköy ($39^{\circ}59'36''N$, $39^{\circ}23'6''E$), 1800 m. 2♀♀; Kırklareli-Dereköy ($41^{\circ}93'45''N$, $27^{\circ}36'60''E$), 450 m. 06.08.1995, 1♀; Siirt-Merkez ($37^{\circ}55'12''N$, $41^{\circ}56'23''E$), 925 m. 30.07.2009 8♀♀, 4♂♂.

Distribution: Palearctic.

Heterospilus (Heterospilus) tadzhicus Belokobylskij, 1983

Material examined: Siirt-Merkez ($37^{\circ}55'31''N$, $41^{\circ}55'32''E$), 880 m. 01.07.2009, 2♀♀; 22.07.2009, 1♀; (Fıstık) 22.07.2009, 1♀; Zonguldak-Çaycuma-Kayıkcılar ($41^{\circ}25'12''N$, $32^{\circ}04'47''E$), 20 m. 30.06.2001, 1♀.

Distribution: Eastern Palearctic.

Heterospilus (Heterospilus) tauricus Telenga, 1941

Material examined: Adana-Balcalı ($37^{\circ}03'26''N$, $35^{\circ}21'43''E$), 138 m. 15.05.1983, 1♀; Adapazarı-Emirdağ-Mezarlık ($40^{\circ}49'37''N$, $30^{\circ}23'16''E$), 150 m. 12.09.1992, 1♀; -Hendek-Hüseyinşeyh ($37^{\circ}20'26''N$, $28^{\circ}8'25''E$), 240 m. 27.06.2001, 3♀♀; -TİGEM ($40^{\circ}47'19''N$, $30^{\circ}24'21''E$), 62 m. 06.07.1996, 1♀; Ankara-Temelli-Alagöz ($39^{\circ}44'25''N$, $32^{\circ}22'40''E$), 803 m. 10.07.2007, 2♀♀; Aydın-Adanan Menderes Üniversitesi Ziraat Fakültesi ($37^{\circ}50'39''N$, $27^{\circ}50'44''E$), 70 m. 18.09.1995, 1♀; Bartın-Merkez ($41^{\circ}38'4''N$, $32^{\circ}20'15''E$), 10 m. 09.06.2002, 2♀♀; -Çamlık ($41^{\circ}32'45''N$, $32^{\circ}27'18''E$), 60 m. 30.06.2001, 1♀; Edirne-Güllapoğlu ($41^{\circ}38'20''N$, $26^{\circ}36'53''E$), 41 m. 03.10.2001, 1♀; -Haciumur (Gece) ($41^{\circ}43'02''N$, $26^{\circ}47'16''E$), 50 m. 27.06.1991, 1♀; -Suakacağı ($41^{\circ}50'25''N$, $26^{\circ}35'18''E$), 210 m. 29.05.1993, 1♀; Elazığ-Keban-İbolar ($37^{\circ}28'54''N$, $38^{\circ}19'3''E$), 888 m. 16.10.2007, 1♀; Erzincan-İliç-Bağışdaş ($39^{\circ}26'46''N$, $38^{\circ}29'38''E$), 1090 m. 26.08.2008, 1♀; -Kemah-Soğukpınar ($39^{\circ}36'06''N$, $39^{\circ}02'17''E$), 1040 m. 26.08.2008, 1♀; Karabük-Bostanbüyük ($41^{\circ}12'53''N$, $32^{\circ}41'21''E$), 290 m. 28.08.2002, 1♀; -Safranbolu-İnceçay-Sarıçıçek Dağı ($41^{\circ}26'55''N$, $32^{\circ}45'13''E$), 1000 m. 29.06.2001, 1♀; -Bostanbüyük ($41^{\circ}12'53''N$, $32^{\circ}41'21''E$), 290 m. 13.06.2002, 1♀; Kastamonu-Araç-Akgeçit ($41^{\circ}13'13''N$, $33^{\circ}12'25''E$), 400 m. 28.06.2002, 1♀; -Azdavay-Kiraz Dağı ($41^{\circ}42'00''N$, $33^{\circ}23'00''E$), 914 m. 05.08.2001, 1♀; -Azdavay-Mehmetçelebi ($41^{\circ}38'26''N$, $33^{\circ}17'53''E$), 730 m. 05.09.2001, 1♀; -Doğanyurt-Denizbüyük ($41^{\circ}59'31''N$, $33^{\circ}29'33''E$), 200 m. 10.06.2002, 1♀; -Küre-Ersizlerdere ($41^{\circ}50'38''N$, $33^{\circ}43'38''E$), 1560 m. 12.06.2002, 1♀; -Küre-Ersizlerdere-İpsizler ($41^{\circ}50'38''N$, $33^{\circ}43'38''E$), 1910 m. 12.06.2002, 1♀; Kırklareli-İnce-Paşayeri ($41^{\circ}40'40''N$, $27^{\circ}4'8''E$), 150 m. 02.08.2000, 2♀♀; Muğla-Yatağan-Dipsiz ($37^{\circ}25'00''N$, $28^{\circ}06'00''E$), 376 m. 31.07.1997, 1♀; Nevşehir-Avanos-Saruhan ($38^{\circ}42'47''N$, $34^{\circ}54'47''E$), 958 m. 06.06.2007, 1♀; Samsun-Havsa, Mismilağaç ($40^{\circ}59'51''N$, $35^{\circ}41'49''E$), 610 m. 02.07.2003, 1♀; -Vezirköprü-Erduaşı ($41^{\circ}21'27''N$, $35^{\circ}03'06''E$), 110 m. 02.07.2003, 1♀; -Vezirköprü-İncesu ($41^{\circ}9'46''N$, $35^{\circ}19'31''E$), 240 m. 28.08.2003, 2♀♀; Sinop-A-B.Zaviye ($41^{\circ}52'52''N$, $34^{\circ}32'8''E$), 16 m. 06.09.2001, 1♀; -Yeniçam ($41^{\circ}96'5''N$, $34^{\circ}85'6''E$), 30 m. 02.07.2001, 1♀; Sivas-Merkez ($39^{\circ}44'24''N$, $37^{\circ}00'35''E$), 1273 m. 17.07.1990, 1♀; Tekirdağ-Kınıklar ($40^{\circ}53'8''N$, $26^{\circ}54'49''E$), 110 m.

26.06.2003, 1♀; -Muratlı (41°10'26"N, 27°30'31"E), 60 m.
 28.08.1992, 1♀; -Naipköy (40°50'49"N, 27°26'48"E), 20 m.
 24.06.2003, 1♀; -Şarköy (40°36'48"N, 27°6'48"E), 120 m.
 25.06.2003, 1♀; -Yazırköy (40°54'26"N, 27°28'7"E), 60 m.
 23.08.2000, 1♀; Tokat-Pazar-Balıca (40°13'29"N, 36°17'34"E), 1020 m. 07.07.2003, 1♀; -Taşlıçiftlik (40°19'19"N, 36°28'13"E), 550 m. 06.07.2003, 1♀; Zonguldak-Çukurören (41°27'23"N, 31°47'55"E), 70 m. 29.06.2001, 1♀.

Distribution: Palearctic.

Host: Coleoptera, Mordellidae: *Mordellistena (Mordellistena) pentas* Mulsant 1856.

3.12. Genus *Hormius* Nees, 1819

Hormius moniliatus (Nees, 1811)

Material examined: Adiyaman-Merkez (37°45'51"N, 38°16'34"E), 676 m. 25.05.2009, 5♀♂; -Besni-Sofraz (37°38'16"N, 37°54'21"E), 696 m. 17.10.2005, 1♀; -Gölbaşı-Çöplü (37°46'9"N, 37°42'4"E), 1030 m. 15.09.2004, 1♂; Aksaray-Ihlara (38°15'27"N, 34°17'43"E), 1212 m. 11.09.2006, 1♂; -Ihlara-Yaprakhisar (38°13'18"N, 34°17'14"E), 1354 m. 11.09.2006, 2♂♂; Amasya-Taşova-Keçiyatağı (40°48'49"N, 36°09'41"E), 1066 m. 08.07.2003, 1♀; Ankara-Elmadağ (39°55'15"N, 33°13'50"E), 1165 m. 16.09.2006, 1♀; -Haymana-Balçıkhisar (39°14'46"N, 32°45'00"E), 1080 m. 10.07.2007, 1♀; -Kalecik-Elmapınar (40°16'47"N, 33°28'04"E), 912 m. 28.05.2007, 1♂; -Kazan-Aydın (40°09'56"N, 32°40'09"E), 872 m. 17.09.2006, 6♀♀; -Kızılcahamam-Pazar (40°20'01"N, 32°42'04"E), 959 m. 17.09.2006, 1♀, 2♂♂; -Yenikent-İlyakut (40°04'19"N, 32°28'16"E), 969 m. 08.06.2007, 1♀; Ardahan-Çamlıçatak (41°07'18"N, 42°50'19"E), 1850 m. 15.09.2012, 1♀; -Çıldır-Eskibeyrehatun Köyü (41°07'47"N, 42°54'32"E), 1950 m. 22.08.2013, 1♀; Artvin-Yusufeli-Sarıgöl (40°57'22"N, 41°29'47"E), 842 m. 04.07.2004, 1♀; Aydın-Kuyucak-Pamukören (37°55'23"N, 28°32'30"E), 140 m. 19.06.1996, 1♀; Balıkesir-Merkez-Ovaköy (39°38'00"N, 28°00'00"E), 95 m. 17.09.1992, 2♀♀; Bartın-Kuruçasıle (41°50'35"N, 32°42'49"E), 129 m. 27.05.2007, 1♀; Bilecik-Başköy (40°5'32"N, 30°0'22"E), 62 m. 20.09.1992, 1♂; Çanakkale-Küçükkyuyu (39°32'57"N, 26°36'34"E), 142 m. 04.11.2004, 1♀; Çankırı-Korgun (40°44'00"N, 33°31'25"E), 898 m. 28.05.2007, 1♀; Denizli-Buldan-Kadıköy (38°9'43"N, 28°45'32"E), 580 m. 19.09.1996, 2♀♀; Diyarbakır-Çermik-Kızılızbük köprüsü (38°8'13"N, 39°27'3"E), 845 m. 04.07.2010, 1♀; Devegeçidi (38°06'34"N, 40°12'46"E), 602 m. 04.07.2010, 9, 1♀; -Ergani (38°15'59"N, 39°45'59"E), 872 m. 26.06.2007, 4♀♀; -Ergani-Pınarkaya (38°14'46"N, 39°41'10"E), 935 m. 21.06.2007, 1♂; Düzce-Akçakoca-Dadaloğlu (41°03'18"N, 31°11'35"E), 96 m. 26.05.2007, 1♀; Edirne-Trakya Üniv.-Güllapoğlu yerleşkesi (41°38'20"N, 26°36'53"E), 41 m. 25.04.2001, I.Tuzağı, 1♂; Elazığ-Aydınlar (38°48'04"N, 38°53'40"E), 985 m. 16.07.2007, 1♀; -Baskılı-Canbeyler

(38°33'24"N, 38°49'00"E), 1116 m. 03.06.2007, 2♂; -Baskılı-Hacımustafa (38°38'34"N, 38°51'07"E), 1436 m. 16.07.2007, 5♀♀; -Baskılı-Yalındam (38°34'26"N, 38°43'55"E), 1400 m. 10.06.2008, 1♂; -Karaçavuş (38°40'36"N, 38°56'08"E), 1424 m. 12.06.2008, 2♂♂; -Keban-Aşgören (38°47'38"N, 38°44'30"E), 780 m. 04.06.2007, 3♀♀; -Keban-İbolar (38°49'03"N, 38°52'16"E), 888 m. 16.07.2007, 3♀♀, 1♂; -Keban-Ulupınar (38°44'59"N, 38°51'57"E), 1229 m. 11.06.2008, 1♀; -Kovancılar-Pinartepe (38°42'37"N, 39°53'06"E), 916 m. 29.08.2008, 1♀; Erzincan-Eğin-Yuva Köyü (39°15'12"N, 38°30'10"E), 830 m. 26.08.2008, 1♀; -Kemah-Soğukpinar (39°36'06"N, 39°02'17"E), 1040 m. 26.08.2008, 3♀♀; -Refahiye-Salur (39°55'37"N, 38°31'37"E), 1610 m. 22.08.2008, 1♂; -Üzümlü-Demirpinar (39°37'38"N, 39°47'34"E), 1200 m. 24.08.2008, 6♀♀; -Üzümlü-Geyikli (39°43'36"N, 39°40'00"E), 1391 m. 23.08.2008, 2♀♀; -Aşkale (39°55'17"N, 40°42'24"E), 1670 m. 18.8.2013, 1♂; Erzurum-Aziziye-Eskipolat (40°04'08"N, 40°56'00"E), 1860 m. 16.08.2013, 1♂; -Horasan-Orman İşletme Müd. (40°02'06"N, 42°10'41"E), 1585 m. 25.08.2013, 1♂; -İspir-Çamlıkaya (40°37'49"N, 41°10'52"E), 950 m. 16.08.2013, 2♀♀, 1♂; -İspir-Köprüköy (40°25'36"N, 40°58'27"E), 1465 m. 11.09.2012, 2♂; -Oltu-Güzelsoy (40°37'19"N, 42°1'2"E), 1605 m. 12.09.2012, 1♂; -Olur (39°54'35"N, 41°16'32"E), 1853 m. 12.09.2012, 1♀, 1♂, 1♀; -Palandöken Kayak Merk. (39°50'57"N, 41°16'59"E), 2455 m. 18.08.2013, 1♀; -Şenkaya-Değirmenlidere köyü (40°43'53"N, 42°31'09"E), 1820 m. 20.08.2013, 1♂; -Şenkaya-Gazialan (40°33'59"N, 42°20'59"E), 915 m. 13.09.2012, 1♀; Eskişehir'e 26 km kala (39°77'93"N, 30°50'79"E), 793 m. 04.09.2006, 2♀♀, 1♂; -Alpu-Sündiken dağları (39°58'36"N, 31°06'22"E), 1469 m. 05.09.2006, 1♂; -Yörükhirka (39°36'00"N, 30°25'12"E), 862 m. 08.07.2007, 2♀♀; Gümüşhane-Şiran-Arıtaş (40°09'43"N, 39°01'40"E), 1300 m. 02.07.2004, 1♀; İğdır-Tuzluca-Gaziler (40°06'36"N, 43°25'31"E), 1035 m. 21.0.2012, 1♀, 1♀; -Tuzluca-Köprübaşı (40°3'6"N, 43°43'31"E), 1035 m. 20.09.2012, 1♀, 1♂, 2♂♂; Isparta-Eğirdir (38°1'25"N, 30°52'22"E), 969 m. 09.08.1980, 1♂; -Keçiborlu (37°56'53"N, 30°18'21"E), 866 m. 31.05.2002, 1♂; İzmit-Sapanca-İst. Üniv. Tesisi (40°11'40"N, 30°31'13"E), 222 m. 16.08.1994, 1♀; Karabük-Safranbolu-İnceçay-Sarıçık Dağı (41°26'55"N, 32°45'13"E), 1000 m. 29.06.2001, 1♀; Kars-Arpaçay-Çanaklısu (41°00'51"N, 43°18'14"E), 1975 m. 22.08.2013, 1♀; -Kağızman-Aydinkavak (40°09'11"N, 43°19'14"E), 1115 m. 20.09.2012, 2♀; 1♂; -Kağızman-Yalnızça (40°13'49"N, 42°59'05"E), 1405 m. 23.08.2013, 1♀, 1♂; Sarıkamış Orm. (40°16'50"N, 42°39'01"E), 2045 m. 24.08.2013, 1♂; -Sarıkamış-Bayraktepe (40°17'28"N, 42°36'00"E), 2644 m. 19.09.2012, 1♂; 24.06.2013, 1♂; -Sarıkamış-Horasan Yolu (40°19'54"N, 42°35'42"E), 2102 m. 21.09.2012, 1♂; -Sarıkamış-Karakurt (40°10'03"N,

42°36'36"E), 1475 m. 24.08.2013, 1♀; -Sarıkamış-Söğütlüçesme (40°16'58"N, 42°34'18"E), 2350 m. 24.06.2013, 1♂; -Susuz-İncesu (40°06'36"N, 43°25'31"E), 1057 m. 17.09.2012, 1♀; Kayseri-Bağpınar (38°49'12"N, 35°37'56"E), 1097 m. 14.09.2006, 1♂; -Bünyan-Ekrek (38°39'42"N, 36°03'30"E), 1424 m. 12.07.2007, 1♂; -Bünyan-Erkek (38°39'42"N, 36°03'30"E), 1424 m. 12.07.2007, 1♀; 22.06.2009, 1♂; -Bünyan (38°50'53"N, 35°51'36"E), 1350 m. 14.09.2006, 3♂♂; -Gömeç (38°52'17"N, 35°39'43"E), 1094 m. 14.09.2006, 3♂♂; -İncesu-Örenşehir (38°41'28"N, 35°14'37"E), 1058 m. 14.09.2006, 1♂; -Pınarbaşı-Aşağıkızılıçevik (38°51'56"N, 36°33'28"E), 1520 m. 13.07.2007, 1♂; -Yahalı-M.Beyli (38°5'59"N, 35°21'38"E), 1210 m. 15.09.2006, 1♀, 1♂; -Yeşilhisar-Kuşcu (38°15'01"N, 35°08'56"E), 1056 m. 15.09.2006, 1♀, 1♂; Kırklareli-Demirköy-Boztaş (41°54'55"N, 27°38'49"E), 350 m. 06.07.1997, 1♂; Kırşehir-Özbağ (39°14'32"N, 34°07'43"E), 1036 m. 16.09.2006, 1♂; -Özbağ (39°14'32"N, 34°07'43"E), 1036 m. 16.09.2006, 2♀♀; -Özbağ (39°14'32"N, 34°07'43"E), 1036 m. 16.09.2006, 1♀; -Özbağ (39°14'32"N, 34°07'43"E), 1036 m. 16.09.2006, 2♀♀, 3♂; Konya-Akşehir-Ulupınar (38°26'24"N, 31°19'51"E), 978 m. 07.9.2006, 2♂♂; -Altınapa Barajı (37°52'46"N, 32°19'44"E), 1280 m. 09.08.2006, 2♂♂; -Beyşehir-Çukurağıl (37°53'21"N, 31°54'20"E), 1228 m. 09.09.2006, 1♀; -Beyşehir-Çukurağıl (37°53'21"N, 31°54'20"E), 1228 m. 09.09.2006, 1♀, 1♂; -Seydişehir (37°28'20"N, 31°49'30"E), 1130 m. 09.09.2006, 2♂; Kütahya-Çavdaralan (39°25'12"N, 29°58'48"E), 1400 m. 29.07.1997, 1♂; Malatya-Akçadağ-Darıca (38°20'39"N, 37°40'22"E), 1540 m. 05.06.2007, 2♂♂; -Akçadağ-Kürecik (38°20'49"N, 37°46'10"E), 1749 m. 23.06.2009, 1♂; -Çiftlik (38°20'01"N, 38°27'31"E), 909 m. 05.06.2007, 1♀; -Darende (38°33'10"N, 37°30'13"E), 1033 m. 13.07.2007, 4♀♀; -Darende-Gümüşdere (38°35'52"N, 37°27'50"E), 1377 m. 30.08.2008, 1♀; -Doğanşehir-Reşadiye (37°59'01"N, 38°00'50"E), 1346 m. 14.07.2007, 1♀; -Doğanşehir-Takaz (38°00'35"N, 37°59'28"E), 1286 m. 08.06.2008, 1♀; -Hekimhan (38°49'44"N, 37°55'00"E), 1094 m. 09.06.2008, 1♀; -Yazihan-Surur (38°31'56"N, 38°15'13"E), 710 m. 08.06.2008, 1♀; Muğla-Milas-Konucuk Köyü (37°17'55"N, 27°54'42"E), 650 m. 24.06.1998, 1♀; Nevşehir-Avanos-Saruhan (38°42'47"N, 34°54'47"E), 958 m. 13.09.2006, 2♂♂; -Avanos-Saruhan (38°42'47"N, 34°54'47"E), 958 m. 13.09.2006, 1♂; -Çalış (38°59'31"N, 34°51'46"E), 1233 m. 05.06.2008, 2♀♀; -Gülşehir-Gülyeri (38°44'55"N, 34°40'27"E), 910 m. 12.09.2006, 4♀♀, 5♂♂; -Gülşehir-Gümüşkent (38°51'12"N, 34°33'44"E), 1286 m. 07.06.2007, 1♂; -Ürgüp (38°36'33"N, 34°54'46"E), 1082 m. 13.09.2006, 1♂; Ordu-Ünye-İnkur (41°2'19"N, 37°12'56"E), 380 m. 05.07.2003, 1♀; Samsun-Vezirköprü-İncesu (41°9'6"N,

35°19'31"E), 240 m. 28.08.2003, 1♀; Siirt-Merkez (37°55'31"N, 41°55'32"E), 880 m. 28.08.2009, 4♀♀; Sivas-Gürün-Gökpinar (38°39'26"N, 37°18'15"E), 1472 m. 13.07.2007, 1♂; -Gürün-İncesu (38°45'36"N, 37°6'30"E), 1735 m. 13.07.2007, 1♀; -Kangal-Yeşildere (39°08'33"N, 37°30'00"E), 1600 m. 01.06.2007, 1♀; -Zara-Ağalıkçay (39°32'11"N, 37°45'39"E), 1427 m. 31.05.2007, 1♂; Tokat-Niksar-Çamiçi (40°35'25"N, 36°57'12"E), 1120 m. 05.07.2003, 1♀; -Pazar-Balıca (40°13'29"N, 36°17'34"E), 1020 m. 07.07.2003, 2♀♀; -Reşadiye (41°21'46"N, 37°25'48"E), 520 m. 28.08.2004, 1♀; Tunceli Merkez-Aktuluk (39°3'38"N, 39°25'36"E), 941 m. 27.08.2008, 13♀♀, 2♂♂; -Pertek-Beydamı (38°57'07"N, 39°27'59"E), 1267 m. 28.08.2008, 1♀; -Pülümür-Hiver (39°29'12"N, 39°53'56"E), 1831 m. 27.08.2008, 1♀; -Pülümür-Kangallı (39°29'0"N, 39°54'10"E), 1343 m. 27.08.2008, 1♀; Yozgat-Akdağmadeni-Gökdere (39°41'38"N, 35°42'02"E), 1194 m. 30.07.2007, 1♀; -Mahmatlı (39°41'51"N, 35°22'45"E), 1080 m. 21.08.2008, 1♀; -Şefaatli-Karanlıkdere (39°34'28"N, 34°38'48"E), 840 m. 29.05.2007, 1♀.

Distribution: Holarctic, Nearctic, Oceanic, Oriental.

Hosts: Lepidoptera, Crambidae: *Achyra nudalis* (Hübner, 1796); *Hellula undalis* (Fabricius, 1781); *Paratalanta hyalinalis* (Hübner, 1796); *Pyrausta aurata* (Scopoli, 1763); *P.purpuralis* (Linnaeus, 1758); *P.sanguinalis* (Linnaeus, 1767). Coleophoridae: *Coleophora trifariella* (Zeller, 1849). Elachistidae: *Agonopterix assimilella* (Treitschke, 1832). Gelechiidae: *Dichomeris marginella* (Fabricius, 1781). Oecophoridae: *Agonopterix adspersella* (Kollar, 1832); *Depressaria pulcherrimella* (Stainton, 1849); *Pexicopia malvella* (Hübner, 1805). Scythrididae: *Scythris inspersella* (Hübner 1817). Tortricidae: *Archips crataegana* (Hübner, 1799).

Tortricidae: *Pandemis corylana* (Fabricius, 1794).

Hormius similis Szépligeti, 1896

Material examined: Adana-Feke-Efkeli (37°49'11.712"N, 35°54'44.9388"E), 279 m. 02.10.1979, 1♀; Amasya-Taşova-Keçiyatağı (40°48'49"N, 36°09'41"E), 1066 m. 08.07.2003, 1♀; Balıkesir-Burhaniye-Pelitköy (39°26'36.258"N, 26°54'4.824"E), 50 m. 22.06.1998, 1♀; Kırklareli-İğneada (41°53'3.3288"N, 27°59'7.1484"E), 10 m. 27.07.1986, 1♀; -Demirköy-Boztaş (41°54'55.7604"N, 27°38'49.9524"E), 350 m. 06.07.1997, 1♀; Tokat-Kemalpaşa (40°21'50"N, 36°30'21"E), 665 m. 30.06.2004, 1♀.

Distribution: Palearctic, Oriental.

3.13. Genus *Leluthia* Cameron, 1887

***Leluthia (Euhecabolodes) asiatica* (Tobias, 1980)**

Material examined: Mardin-Derik-Zeytin (37°21'56"N, 40°16'10"E), 620 m. 15.05.2011, 1♀.

Distribution: Eastern Palearctic.

Host: Coleoptera, Curculionidae: *Scolytus schevyrewi* Semenov, 1902.

Leluthia (Euhecabolodes) rugulosolyti (Fischer, 1962)

Material examined: Ankara (40°1'15"N, 32°49'51"E), 870 m. 08.08.2010, 1♀; Diyarbakır-Ergani-Şeftali (8°15'59"N, 39°45'59"E), 872 m. 10.06.1998, 1♀; Elazığ-Kayısı (38°40'51"N, 39°13'35"E), 1043 m. 10.06.1998, 12♀♂; Diyarbakır-Ergani (38°15'59"N, 39°45'59"E), 872 m. 02.06.1998, 1♀; Malatya-Tekevler (38°21'18"N, 38°18'34"E), 977 m. 15.08.1997, 2♀♂.

Distribution: Eastern Palearctic.

Hosts: Coleoptera, Curculionidae: *Phloeosinus bicolor* (Brulle, 1832); *Scolytus rugulosus* (Mueller, 1818).

Leluthia (Euhecabolodes) transcaucasica (Tobias, 1976)

Material examined: Siirt-Merkez (37°55'31"N, 41°55'32"E), 880 m. 30.07.2009, 1♀.

Distribution: Palearctic.

Hosts: Coleoptera, Curculionidae: *Phloeosinus bicolor* (Brulle, 1832); *Scolytus butovitschi* Eggers, 1942c; *S. japonicus* Chapius, 1869.

Leluthia (Leluthia) paradoxa (Picard, 1938)

Material examined: Tokat-Çamağzı (40°24'44"N, 36°41'08"E), 692 m. 01.07.2004, 1♂.

Distribution: Western Palearctic.

Hosts: Coleoptera, Curculionidae: *Pissodes piceae* (Illiger, 1807); *Trypodendron signatum* (Fabricius, 1787). Scolytidae: *Scolytus multistriatus* (Marsham, 1802).

3.14. Genus *Oncophanes* Förster, 1862*Oncophanes (Oncophanes) minutus* (Wesmael, 1838)

Material examined: Artvin-Borçka-Cankurtaran (41°24'01"N, 41°31'10"E), 595 m. 30.06.2005, 1♀; Bolu-Aladağ-Kökez (40°43'50"N, 31°36'34"E), 1000 m. 27.08.2002, 1♀; İstanbul-Bahçeköy-Bilezikçi Çiftliği (41°11'33"N, 28°59'53"E), 30 m. 24.06.1993, 1♀; Kırklareli-Dereköy (41°56'4"N, 27°21'57"E), 500 m. 17.07.2001, 1♀; Samsun-Salıpazarı-Derbentaltı (41°4'49"N, 36°49'35"E), 970 m. 03.07.2003, 1♀.

Distribution: Palearctic.

3.15. Genus *Ontsira* Cameron, 1900*Ontsira antica* (Wollaston, 1858)

Material examined: Artvin-Varlık (41°13'16"N, 41°52'02"E), 673 m. 03.09.2004, 1♀; Edirne-Binevler (41°40'12"N, 26°33'02"E), 41 m. 07.06.1986, 1♀; 07.06.1986, 1♀; 09.06.1991, 1♀; Erzurum-Olur (39°54'35"N, 41°16'32"E), 1853 m. 12.09.2012, 1♀.

Distribution: Holarctic, Oceanic.

Hosts: Coleoptera, Anobiidae: *Ptilinus pectinicornis* (Linnaeus, 1758). Buprestidae: *Agrilus viridis* (Linnaeus, 1758); *Chrysobothris (Chrysobothris) igniventris* Reitter, 1895. Cerambycidae: *Clytus arietis* (Linnaeus, 1758); *Ernobius mollis* (Linnaeus, 1758); *Exocentrus punctipennis* Mulsant & Guillebeau, 1856; *Phymatodes testaceus* (Linné, 1758); *Plagionotus arcuatus* (Linnaeus, 1758); *Plagionotus floralis* (Pallas, 1773); *Pogonoherus*

hispidulus (Piller & Mitterpacher, 1783); *Pyrrhidium sanguineum* (Linnaeus, 1758). Curculionidae: *Blastophagus piniperda* (Linnaeus, 1758); *Magdalisch (Magdalisch) frontalis* (Gyllenhal 1827). Eucnemidae: *Melasis buprestoides* (Linnaeus, 1761). Scolytinae: *Ips acuminatus* (Gyllenhal 1827); *Scolytus pygmaeus* (Fabricius, 1787); *S. rugulosus* (Müller, 1818).

Ontsira imperator (Haliday, 1836)

Material examined: Bolu-Çaydurt (40°44'22"N, 31°36'41"E), 720 m. 28.06.2001, 1♂; Isparta-Merkez (37°45'53"N, 30°33'23"E), 1031 m. 15.06.2003, 1♀, 1♂; Kastamonu-Çatören (41°08'00"N, 33°48'00"E), 950 m. 30.08.2002, 1♀; Kırklareli-Demirköy-Karacadağ (41°57'16"N, 27°40'37"E), 285 m. 03.06.1993, 1♀; Nevşehir-Gülhisar-Açıksaray (38°44'16"N, 34°39'55"E), 1000 m. 12.09.2006, 1♀.

Distribution: Holarctic, Oriental.

Hosts: Coleoptera, Buprestidae: *Buprestis strigosa* (Gebler, 1830). Bostrichidae: *Bostrichus capucinus* (Linnaeus, 1758). Depressariidae: *Dicerea berolinensis* (Herbst, 1779). Cerambycidae: *Acanthocinus aedilis* (Linnaeus, 1758); *A. griseus* (Fabricius, 1792); *Anisarthron barbipes* (Schrank, 1781); *Hylotrupes bajulus* (Linnaeus, 1758); *Leiopus nebulosus* (Linnaeus, 1758); *Leptura rubra* (Linnaeus, 1758); *Pogonoherus hispidus* (Linnaeus, 1758); *P. fasciculatus* (De Geer, 1775); *Rhagium bifasciatum* (Fabricius 1775); *Scotylus scotylus* (Fabricius 1775); *Stromatium fulvum* (Linsley & Chemsak, 1997); *Tetropium castaneum* (Linnaeus, 1758). Curculionidae: *Blastophagus piniperda* (Linnaeus 1758). Scolitidae: *Blastophagus minor* (Hartig, 1834).

3.16. Genus *Pachystigmus* Hellén, 1927*Pachystigmus nitidulus* Hellén, 1927

Material examined: Aksaray-Ihlara-Yapraklıhisar (38°13'18"N, 34°17'14"E), 1354 m. 11.09.2006, 1♀, 1♂; Eskişehir-Anadolu Üniversitesi Hatıra Ormanı (39°43'09"N, 30°40'37"E), 907 m. 10.07.2007, 2♀♂; Sivrihisar-Babatat (39°31'42"N, 31°37'36"E), 868 m. 10.07.2007, 4♀♂; Malatya-Doğanşehir-Kemaliye (37°16'45"N, 38°0'17"E), 1345 m. 02.06.2007, 1♂.

Distribution: Western Palearctic.

3.17. Genus *Pambolus* Haliday, 1836**Pambolus (Phaenodus) pallipes* (Förster, 1862)

Material examined: Nevşehir-Amanos-Saruhan (38°42'47"N, 34°54'47"E), 958 m. 13.09.2006, 1♀; Rize-Ardeşen-Duygulu (47°07'02"N, 41°02'45"E), 132 m. 27.06.2005, 1♂.

Distribution: Palearctic.

3.18. Genus *Polystenus* Förster, 1862*Polystenus rugosus* Förster, 1862

Material examined: Nevşehir-Avanos-Saruhan (38°42'47"N, 34°54'47"E), 958 m. 13.09.2006, 1♀.

Distribution: Palearctic, Oriental.

Hosts: Coleoptera, Buprestidae: *Agrilus* sp.; *Agrilus angustulus* (Illiger, 1803); *A. auricollis* (Kiesenwetter, 1857); *A. sulcicollis* Lacordaire, 1835; *A. viridis* (Linnaeus, 1758); *Anthaxia manca* (Linnaeus, 1767); *Coraebus florentinus* (Herbst, 1801) syn. *C. bifasciatus* Olivier; *Sinoxylon sexdentatum*; (Olivier, 1790).

3.19. Genus *Pseudohormius* Tobias & Alexeev, 1973

Pseudohormius turkmenus Tobias & Alexeev, 1973

Material examined: Elazığ-Baskıl-Yalındam (38°34'26"N, 38°43'55"E), 1400 m. 10.06.2008, 2♀♂; 1♂; Erzincan-Refahiye-Salur (39°55'37"N, 38°31'37"E), 1610 m. 22.08.2008, 1♀; Eskişehir-Kaymaz (39°50'57"N, 31°10'06"E), 449 m. 10.07.2007, 1♀; Kars-Sarıkamış-Erzurum Yolu 10 km (40°15'03"N, 42°39'11"E), 2016 m. 23.06.2013, 1♂; Kayseri-Gömeç (38°52'17"N, 35°39'43"E), 1094 m. 14.09.2006, 1♂; Malatya-Darende-Gümüşdere (38°35'52"N, 37°27'50"E), 1377 m. 30.08.2008, 1♀; -Doğanşehir-Takaz (38°00'35"N, 37°59'28"E), 1286 m. 14.07.2007, 1♂; Sivas-Gürün-İncesu (38°45'36"N, 37°6'30"E), 1735 m. 13.07.2007, 1♀; Tunceli-Merkez-Aktuluk (39°02'29"N, 39°30'25"E), 941 m. 27.08.2008, 1♀; Yozgat-Şefaatlı-Karanlıkdere (39°34'28"N, 34°38'48"E), 840 m. 29.05.2007, 1♂.

Distribution: Palearctic.

Host: Lepidoptera, Buccalatricidae: *Bucculatrix crataegi* Zeller, 1839.

3.20. Genus *Rhaconotus* Ruthe, 1854

Rhaconotus (Rhaconotus) aciculatus Ruthe, 1854

Material examined: Amasya-Yıldızköy (40°34'14"N, 35°42'54"E), 420 m. 23.06.2004, 2♀♀; Ankara-Temelli-Alagöz (39°45'18"N, 32°26'42"E), 803 m. 10.07.2007, 1♀; Diyarbakır-Devegeçidi barj. Yol ayr. (38°06'34"N, 40°12'46"E), 602 m. 04.07.2010, 3♀♀; Edirne-Havsa-Köseömer (41°35'14"N, 26°54'15"E), 115 m. 30.08.1992, 1♀; Erzincan-Üzümlü-Demirpinar (39°37'38"N, 39°47'34"E), 1200 m. 24.08.2008, 1♀; Erzurum-İspir-Ardıçı (40°33'59"N, 41°14'0"E), 1560 m. 19.06.2012, 1♀; İğdır-Karakoyunlu-Taşburun (39°58'55"N, 44°12'03"E), 839 m. 26.06.2012, 2♀♀; İğdır-Tigem (39°47'10"N, 44°37'05"E), 850 m. 26.06.2012, 8♀♀; İğdır-Tuzluca-Gaziler (40°06'36"N, 43°25'31"E), 1035 m. 27.0.2012, 1♀; Malatya-Hekimhan-Yazılı (38°42'46"N, 38°05'06"E), 1176 m. 09.08.2008, 1♀; Muğla-Yatağan-Dipsiz (37°25'00"N, 28°06'00"E), 376 m. 31.07.1997, 1♀; Sivas-Kangal-Yeşildere (39°08'33"N, 37°30'00"E), 1600 m. 01.06.2007, 2♀♀; Tekirdağ-Yazırköy (40°54'26"N, 27°28'7"E), 60 m. 23.08.2000, 2♀♀.

Distribution: Palearctic, Oriental.

Hosts: Coleoptera, Buprestidae: *Agrilus viridis* (Linnaeus, 1758); *Anthaxia lgockii* Obenberger, 1917; *Caryedon* sp. Chrysomelidae: *Caryedon serratus* (Olivier, 1790); *Lixus Glaucopteron lukjanovitschi* Putschkov, 1975.

Rhaconotus (Rhaconotus) asiaticus Belokobylskij, 1990

Material examined: Afyon-Emirdağ-Kırkpınar (39°12'24"N, 31°01'56"E), 911 m. 07.07.2006, 1♂; Amasya-Merzifon-Tavşan D.-Esenköy Y. (41°00'37"N, 35°17'23"E), Ankara-Temelli-Alagöz (39°45'18"N, 32°26'42"E), 803 m. 10.07.2007, 1♀; Antalya-*Prosopis forcta* (36°12'0"N, 36°10'34"E), 86 m. 10.10.2001, 1♀; Artvin-Dereci (41°28'52"N, 41°31'34"E), 1018 m. 01.07.2005, 1♀; Aydın-Germencik-Morali (37°53'23"N, 27°30'16"E), 100 m. 24.06.1998, 1♀; Bartın-Kuruca-Şile-Danişment (41°49'37"N, 32°41'23"E), 126 m. 09.06.2002, 1♀; Bayburt-Merkez (40°14'51"N, 40°13'46"E), 1562 m. 07.08.2005, 1♀; Denizli-Buldan-Kadıköy (38°9'43"N, 28°45'32"E), 580 m. 19.09.1996, 1♀; Diyarbakır (37°54'51"N, 40°13'50"E), 641 m. 05.05.2006, 1♀; Edirne-Enez-Sultaniçi (40°37'37"N, 26°9'29"E), 20 m. 22.08.1991, 1♀; -Uzunköprü-Kircasalih (41°23'57"N, 26°48'36"E), 100 m. 01.08.1992, 1♀; Erzincan-Üzümlü (39°41'04"N, 39°39'32"E), 1162 m. 23.08.2008, 1♀; -Kemah-Soğukpinar (39°36'06"N, 39°02'17"E), 1040 m. 26.08.2008, 1♀; -Tercan-Elmalı (39°34'17"N, 40°08'35"E), 1350 m. 24.08.2008, 1♂; -Üzümlü-Geyikli (39°43'36"N, 39°40'00"E), 1391 m. 15.06.2008, 1♀; Erzurum (39°54'15"N, 41°16'4"E), 1889 m. 29.07.1994, 1♀; Eskişehir-Türkmendağı-Kavacık (39°31'59"N, 30°20'28"E), 1230 m. 08.07.2007, 1♀; Gökçeada-TİGEM (40°11'49"N, 25°55'4"E), 35 m. 06.07.1996, 1♀; Kastamonu-Azday-Mehmetçelebi (41°38'26"N, 33°17'53"E), 730 m. 05.09.2001, 1♀; Kayseri-Bünyan (38°50'53"N, 35°51'36"E), 1350 m. 12.07.2007, 1♀; Malatya-Doğanşehir-Takaz (38°00'35"N, 37°59'28"E), 1286 m. 14.07.2007, 1♀; Nevşehir-Hacıbektaş (38°57'43"N, 34°33'29"E), 1255 m. 07.06.2007, 1♀; Sinop-Boyabat-Çangal Dağı-Kozcağız (41°27'36"N, 34°46'48"E), 333 m. 1♂; Sivas-Zara (39°53'42"N, 37°45'11"E), 1010 m. 31.05.2007, 1♀.

Distribution: Eastern Palearctic.

Host: Coleoptera, Curculioniae: *Lixus (Eulixus) kiritshenkoi* Ter-Minasyan 1985.

Rhaconotus (Rhaconotus) elegans (Förster, 1862)

Material examined: Ankara-Yenitermidal (40°07'41"N, 32°59'42"E), 952 m. 27.10.2001, 1♀.

Distribution: Palearctic.

Rhaconotus (Rhaconotus) kerzhneri Belokobylskij, 1985

Material examined: Antalya-*Prosopis forcta* (36°12'0"N, 36°10'34"E), 86 m. October 2001, 1♀; İçel-Erdemli (36°36'05"N, 34°18'1"E), 15 m. 06.07.1983, 1♂.

Distribution: Eastern Palearctic.

Rhaconotus (Rhaconotus) longulus Belokobylskij, 1994

Material examined: Bozcaada-Merkez (39°49'23"N, 26°1'57"E), 8 m. 08.07.1996, 1♀; Çanakkale-Eceabat

($40^{\circ}11'1''N$, $26^{\circ}21'22''E$), 9 m. 15.07.1986, 1♀; İğdır-Karakoyunlu-Taşburun ($39^{\circ}58'55''N$, $44^{\circ}12'03''E$), 839 m. 1♂; Tekirdağ-Işıkalar ($40^{\circ}48'54''N$, $27^{\circ}22'50''E$), 400 m. 14.07.1986, 1♀.

Distribution: Eastern Palearctic.

Rhaconotus (Rhaconotus) pictipennis (Reinhard, 1885)

Material examined: Balıkesir-Burhaniye-Pelitköy ($39^{\circ}26'36''N$, $26^{\circ}54'4''E$), 50 m. 22.06.1998, 1♂; Bartın-Merkez ($41^{\circ}38'4''N$, $32^{\circ}20'15''E$), 10 m. 09.06.2002, 1♀; Elazığ-Kovancılar-Konakalmaz ($38^{\circ}40'44''N$, $39^{\circ}42'59''E$), 1161 m. 29.08.2008, 1♂; Eskişehir(26km kala) ($39^{\circ}46'48''N$, $30^{\circ}30'36''E$), 793 m. 04.09.2006, 1♀; Kastamonu-Araç-Toprakcuma ($41^{\circ}23'56''N$, $33^{\circ}46'46''E$), 350 m. 13.06.2002 1♀; -Hanönü ($41^{\circ}37'33''N$, $34^{\circ}27'43''E$), 450 m. 02.07.2001, 1♀; Tunceli-Merkez-Aktuluk ($39^{\circ}3'38''N$, $39^{\circ}25'36''E$), 941 m. 27.08.2008, 1♀; Zonguldak-Çaycuma-Kayıççilar ($41^{\circ}25'12''N$, $32^{\circ}04'47''E$), 20 m. 30.06.2001, 1♀, 1♂; Zonguldak-Karadeniz Ereğlisi ($41^{\circ}17'21''N$, $31^{\circ}25'5''E$), 40 m. 08.06.2002, 1♀.

Distribution: Palearctic.

3.21. Genus *Rhoptrocentrus* Marshall, 1897

Rhoptrocentrus piceus Marshall, 1897

Material examined: Amasya-Yıldızköy ($40^{\circ}34'14''N$, $35^{\circ}42'54''E$), 420 m. 23.06.2004, 1♀; Edirne-Trakya Üniv. Merkez Kampüsü-(Zooloji Lab.) ($41^{\circ}38'20''N$, $26^{\circ}36'53''E$), 41 m. 16.10.1989, 1♀.

Distribution: Palearctic.

3.22. Genus *Rhysipolis* Förster, 1862

Rhysipolis decorator (Haliday, 1836)

Material examined: Afyon-Çay ($38^{\circ}35'33''N$, $31^{\circ}03'03''E$), 1004 m. 07.07.2006, 1♀; Aksaray-Ihlara ($38^{\circ}15'27''N$, $34^{\circ}17'43''E$), 1212 m. 11.09.2006, 1♀; Yapraklıhisar ($38^{\circ}13'18''N$, $34^{\circ}17'14''E$), 1354 m. 11.09.2006, 1♀; Amasya-Taşova-Boraboy köyü ($40^{\circ}48'35''N$, $36^{\circ}12'07''E$), 551 m. 08.07.2003, 1♀; Ankara-Kızılcahamam-Özbekler ($40^{\circ}32'32''N$, $32^{\circ}36'29''E$), 970 m. 17.09.2006, 1♀; -Yenikent-Ilyakut ($40^{\circ}04'19''N$, $32^{\circ}28'16''E$), 969 m. 08.06.2007, 1♀; Ardahan-Kent Ormanı ($41^{\circ}6'37''N$, $42^{\circ}42'7''E$), 1810 m. 15.09.2012, 2♀♀; Artvin-Yusufeli-Küplüce ($40^{\circ}56'06''N$, $40^{\circ}30'23''E$), 828 m. 04.09.2004, 1♀; Bayburt-Konurusu ($40^{\circ}23'39''N$, $40^{\circ}14'54''E$), 1522 m. 30.08.2004, 1♀; -Merkez ($40^{\circ}14'51''N$, $40^{\circ}13'46''E$), 1562 m. 07.08.2005, 1♀; Çankırı-Korgun ($40^{\circ}44'00''N$, $33^{\circ}31'25''E$), 898 m. 28.05.2007, 2♀♀; Denizli-Buldan-Kadıköy ($38^{\circ}9'43''N$, $28^{\circ}45'32''E$), 580 m. 19.09.1996, 1♀; Elazığ-Aydınlar ($38^{\circ}48'04''N$, $38^{\circ}53'40''E$), 985 m. 16.07.2007, 1♀; -Baskıl-Hacımustafa ($38^{\circ}38'34''N$, $38^{\circ}51'07''E$), 1436 m. 23.07.1985, 1♀; Erzincan-Üzümlü ($39^{\circ}41'04''N$, $39^{\circ}39'32''E$), 1162 m. 23.08.2008, 4♀♀; -Üzümlü-Geyikli ($39^{\circ}43'36''N$, $39^{\circ}40'00''E$), 1391 m. 23.08.2008, 1♀; Eskişehir-A.Ü-Hatıra Ormanı ($39^{\circ}43'09''N$, $30^{\circ}40'37''E$),

907 m. 10.07.2007, 1♀; -Kaymaz ($39^{\circ}50'57''N$, $31^{\circ}10'06''E$), 449 m. 10.07.2007, 2♀♀; Gümüşhane-Kelkit-Köycük ($40^{\circ}9'1''N$, $39^{\circ}25'19''E$), 1450 m. 08.08.2005, 1♀; Kastamonu-Azdadavay-Mehmetçelebi ($41^{\circ}38'26''N$, $33^{\circ}17'53''E$), 730 m. 05.09.2001, 1♀; -Daday-Ballıdağ ($41^{\circ}28'0''N$, $33^{\circ}25'0''E$), 1750 m. 01.07.2001, 1♀; Kayseri-Bağpınar ($38^{\circ}49'12''N$, $35^{\circ}37'56''E$), 1097 m. 14.09.2006, 8♀♀, 2♂♂; -Gömeç ($38^{\circ}52'17''N$, $35^{\circ}39'43''E$), 1094 m. 14.09.2006, 1♀; -Yemiliha ($38^{\circ}52'02''N$, $35^{\circ}14'32''E$), 992 m. 15.09.2006, 3♀♀, 1♂; -Yeşilhisar-Araplı ($38^{\circ}14'05''N$, $35^{\circ}03'13''E$), 1349 m. 18.07.2007, 3♀♀; Kırıkkale-Karakeçili ($39^{\circ}34'35''N$, $33^{\circ}21'25''E$), 839 m. 11.07.2007, 2♀♀; Kırşehir-Mucur-Kurugöl ($39^{\circ}01'54''N$, $34^{\circ}26'41''E$), 1046 m. 07.06.2007, 1♀; Malatya-Akçadağ-Darıca ($38^{\circ}20'39''N$, $37^{\circ}40'22''E$), 1540 m. 05.06.2007, 1♀; -Çiftlik ($38^{\circ}20'01''N$, $38^{\circ}27'31''E$), 909 m. 18.07.2007, 1♀; Muğla-Yatağan-Dipsiz ($37^{\circ}25'00''N$, $28^{\circ}06'00''E$), 376 m. 31.07.1997, 1♀; Nevşehir-Avanos-Saruhan ($38^{\circ}42'47''N$, $34^{\circ}54'47''E$), 958 m. 13.09.2006, 2♀♀; -Ürgüp ($38^{\circ}36'33''N$, $34^{\circ}54'46''E$), 1082 m. 13.09.2006, 2♀♀, 1♂; Sivas-Kampüs ($39^{\circ}44'51''N$, $37^{\circ}1'43''E$), 1285 m. 24.08.2002, 1♀; Tokat-Erbaa ($40^{\circ}40'09''N$, $36^{\circ}38'06''E$), 234 m. 28.08.2004, 1♀; -Pazar-Balıca ($40^{\circ}13'29''N$, $36^{\circ}17'34''E$), 1020 m. 07.07.2003, 1♀; -Turhal-Doğanlı Çiftliği ($40^{\circ}18'14''N$, $36^{\circ}19'29''E$), 554 m. 30.06.2004, 2♂♂; Trabzon-Akçabat-Akat ($40^{\circ}59'01''N$, $39^{\circ}33'06''E$), 77 m. 04.08.2005, 1♀; Tunceli-Pülümür-Kangalı ($39^{\circ}27'10''N$, $39^{\circ}51'43''E$), 1343 m. 27.08.2008, 1♀.

Distribution: Holarctic.

Hosts: Lepidoptera, Crambidae: *Loxostege sticticalis* (Linnaeus, 1761). Gelechiidae: *Scrobipalpa atriplicella* (Röslerstamm, 1841); *Syncopacma coronillella* (Treitschke, 1833). Gracillariidae: *Acrocercops brongniardella* (Fabricius, 1798); *Caloptilia* sp.; *Caloptilia alchimiella* (Scopoli, 1763); *C. anthobaphes* (Meyrick, 1921); *C. betulincola* (M. Hering, 1928); *C. robustella* Jäckh, 1972; *Choreutis pariana* (Clerck, 1759); *Parornix alta* (Braun, 1925); *P. geminatella* (Packard, 1869); *Phyllonorycter* sp.; *Phyllonorycter harrisella* (Linnaeus, 1761); *P. ostryaefoliella* (Clemens, 1859); *P. propinquella* (Braun, 1908); *P. ulmifoliella* (Hübner, 1817). Lyonetiidae: *Leucoptera malifoliella* (Costa, 1836). Momphidae: *Mompha raschkiella* (Zeller, 1839). Diptera, Anthomyiidae: *Pegomya hyoscyami* (Panzer, 1809). Tephritidae: *Euphranta* (*Euphranta*) *connexa* (Fabricius 1794); *Plioreocepta poeciloptera* (Schränk, 1776); *Tachydromia connexa* Meigen, 1822.

Rhysipolis enukidzei Tobias, 1976

Material examined: Ardahan-Çıldır-Eskibeyrehatun köyü ($41^{\circ}07'7''N$, $42^{\circ}54'32''E$), 1950 m. 22.08.2013, 1♀; Elaçığ-Palu-Seydili ($38^{\circ}41'36''N$, $39^{\circ}52'51''E$), 880 m. 29.08.2008, 1♀; İğdir-Tuzluca ($40^{\circ}02'51''N$, $43^{\circ}42'54''E$), 1038 m. 23.08.2013, 1♀; -Tuzluca-İncesu ($40^{\circ}06'36''N$,

43°25'31"E), 1057 m. 23.06.2013, 2♀♀; Kars-Kağızman-Morpet (40°15'16"N, 42°58'36"E), 1486 m. 23.08.2013, 1♀; -Kağızman-Yalnızağac (40°13'49"N, 42°59'05"E), 1405 m. 23.08.2013, 1♀; -Sarıkamış-Büyükkumru (40°25'14"N, 42°30'52"E), 2102 m. 25.06.2013, 2♀♀; -Sarıkamış-Karakurt (40°10'03"N, 42°36'36"E), 1475 m. 23.06.2013, 1♂.

Distribution: Palearctic.

Rhysipolis hariolator (Haliday, 1836)

Material examined: Aksaray-Ihlara (38°15'27"N, 34°17'43"E), 1212 m. 11.09.2006, 1♀; Ardahan-Çıldır (41°3'28"N, 43°12'41"E), 1920 m. 16.09.2012, 1♀, 1♂; -Göle-Yiğitkonağı köyü (40°55'36"N, 42°35'35"E), 1015 m. 22.06.2012, 1♀, 1♂; -Merkez-Tepeler (41°6'37"N, 42°42'7"E), 1835 m. 22.06.2012, 1♀; -Posof (41°30'30"N, 42°44'03"E), 1445 m. 21.08.2013, 1♀; 14.09.2012, 1♀; Aydın-Sultanhisar-Salavatlı (37°53'34"N, 28°6'17"E), 80 m. 19.09.1996, 1♀; Denizli-Buldan-Kadıköy (38°9'43"N, 28°45'32"E), 580 m. 19.09.1996, 1♂; Erzincan-Kemah-Soğuksınar (39°36'06"N, 39°02'17"E), 1040 m. 26.08.2008, 1♀; Erzurum-Oltu-Orçuk (40°31'30"N, 41°48'23"E), 1270 m. 13.09.2012, 2♂♂; -Olur (39°54'35"N, 41°16'32"E), 1853 m. 12.09.2012, 4♀♀; -Şenkaya-Değirmenlidere (40°43'53"N, 42°31'09"E), 1820 m. 21.06.2012, 1♀; -Şenkaya-Gözalan (40°44'36"N, 42°17'40.0992"E), 1500 m. 13.09.2012, 1♀; -Tortum-Enlem (40°8'00"N, 41°5'58"E), 1515 m. 12.09.2012, 1♂; Eskişehir-Sivrihisar-Babatat (39°31'42"N, 31°37'36"E), 868 m. 10.07.2007, 1♀; Gümüşhane-Kelkit-Yeniköy (40°19'26"N, 39°29'33"E), 1474 m. 29.08.2004, 1♀; İğdır-Tuzluca-Gaziler (40°06'36"N, 43°25'31"E), 1035 m. 27.06.2012, 1♀, 1♂; 21.09.2012, 3♀♀; -Tuzluca-Köprübaşı köyü (40°3'6"N, 43°43'31"E), 1035 m. 25.06.2012, 2♀♀; 20.09.2012, 1♀; Kars-Kağızman (40°9'27"N, 43°7'23"E), 1078 m. 27.06.2012, 1♀; -Kağızman-Aydınkavak (40°09'11"N, 43°19'14"E), 1115 m. 24.08.2013, 3♀♀; 20.09.2012, 3♀♀, 1♂; -Kağızman-Morpet (40°15'16"N, 42°58'36"E), 1486 m. 23.08.2013, 1♀; 20.09.2012, 1♀; -Sarıkamış-Horasan yolu (40°19'54"N, 42°35'42"E), 2102 m. 27.06.2012, 1♀; Kayseri-Bağpınar (38°49'12"N, 35°37'56"E), 1097 m. 14.09.2006, 1♀; -Yemliha (38°52'02"N, 35°14'32"E), 992 m. 15.09.2006, 3♀♀; -Yeşilhisar-Araplı (38°14'05"N, 35°03'13"E), 1349 m. 18.07.2007, 2♂♂; Kırşehir-Daman-Darıözü (39°30'21"N, 33°47'08"E), 902 m. 16.09.2006, 1♀; Konya-Direksarayı (37°24'21"N, 32°36'29"E), 1030 m. 09.09.2006, 1♀; Malatya-Darende-Gümüşdere (38°35'52"N, 37°27'50"E), 1377 m. 30.08.2008, 2♀♀; -Hekimhan (38°49'44"N, 37°55'00"E), 1094 m. 09.06.2008, 2♀♀; Sinop-Yeniçam (41°96'5"N, 34°85'6"E), 30 m. 06.09.2001, 1♀, 1♂; Sivas-Gürün-Suçatı (38°43'17"N, 37°20'45"E), 1266 m. 02.06.2008, 1♀; Tokat-Pazar-Balıca (40°13'29"N, 36°17'34"E), 1020 m. 07.07.2003, 1♀; Tunceli-Pertek-Beydamı (38°56'40"N, 39°28'22"E), 1267

m. 28.08.2008, 2♂♂; -Pülümür-Kangallı (39°27'10"N, 39°51'43"E), 1343 m. 27.08.2008, 1♀.

Distribution: Palearctic.

Hosts: Lepidoptera, Cosmopterigidae: *Cosmopterix zieglerella* Hübner, 1810. Gracillariidae: *Caloptilia alchimiella* (Scopoli, 1763); *C. elongella* (Linnaeus, 1761); *Parornix betulae* (Stainton, 1854); *P. devoniella* (Stainton, 1850); *P. fagivora* (Frey, 1861); *P. petiorella* (Frey, 1863); *P. scotica* (Stainton, 1850); *P. torquillella* (Zeller, 1850); *Phyllonorycter coryli* (Nicelli, 1851); *P. froelichiella* (Zeller, 1839); *P. heegeriella* (Zeller, 1846); *P. kleemannella* (Fabricius, 1781); *P. nicellii* (Stainton, 1851); *P. quercifoliella* (Zeller, 1839).

Rhysipolis meditator (Haliday, 1836)

Material examined: Aksaray-Ihlara (38°15'27"N, 34°17'43"E), 1212 m. 11.09.2006, 1♀; -Ihlara-Yaprakhisar (38°13'18"N, 34°17'14"E), 1354 m. 11.09.2006, 1♂; Amasya-Yolyanı (40°34'22"N, 36°06'38"E), 700 m. 30.06.2004, 1♀; Aydın-Kuyucak-Pamukören (37°55'23"N, 28°32'30"E), 140 m. 19.06.1996, 1♀; -Kuyucak-Pamukören (37°55'23"N, 28°32'30"E), 140 m. 19.09.1996, 1♀, 1♂; -Sultanhisar-Salavatlı (37°53'34"N, 28°6'17"E), 80 m. 19.09.1996, 1♀; Ziraat Fak. (37°50'39.8328"N, 27°50'44.88"E), 47 m. 18.09.1996, 2♀♀; Denizli-Buldan-Kadıköy (38°9'43"N, 28°45'32"E), 580 m. 19.09.1996, 1♂; Edirne-Lalapaşa (41°50'24"N, 26°43'48"E), 560 m. 06.06.1987, 1♀; Elazığ-Baskıl-Hacımustafa Köyü (38°38'34"N, 38°51'07"E), 1436 m. 08.08.1990, 1♀; -Keban-Sağdıçlar (38°46'19"N, 38°50'45"E), 1221 m. 11.06.2008, 1♀; Erzincan-Kemah-Soğuksınar (39°36'06"N, 39°02'17"E), 1040 m. 26.08.2008, 1♂; -Reşadiye-Salur (39°55'37"N, 38°31'37"E), 1610 m. 2.08.2008, 1♀; -Üzümlü-Geyikli (39°43'36"N, 39°40'00"E), 1391 m. 23.08.2008, 1♀; Erzurum-Oltu-Orçuk (40°31'30"N, 41°48'23"E), 1270 m. 13.09.2012, 2♀♀, 2♂♂; -Olur (39°54'35"N, 41°16'32"E), 1853 m. 12.09.2012, 1♀; 1♀; -Şenkaya-Değirmenlidere (40°43'53"N, 42°31'09"E), 1820 m. 21.06.2012, 1♀; -Şenkaya-Gözalan (40°44'36"N, 42°17'40"E), 1500 m. 13.09.2012, 1♀; Eskişehir-Sivrihisar-Babatat (39°31'42"N, 31°37'36"E), 868 m. 10.07.2007, 1♀; -Yörükhirka (39°36'00"N, 30°25'12"E), 862 m. 08.07.2007, 1♀; İğdır-Tuzluca-Köprübaşı (40°3'6.354"N, 43°43'31.476"E), 1035 m. 20.09.2012, 2♀♀; Kars-Boğazköy (40°42'07"N, 43°06'57"E), 1705 m. 17.09.2012, 1♀; -Kağızman-Aydınkavak (40°09'11"N, 43°19'14"E), 1115 m. 20.09.2012, 2♀♀; -Kağızman-Morpet (40°15'16"N, 42°58'36"E), 1486 m. 20.09.2012, 1♀; -Sarıkamış-Bayraktepe (40°17'28"N, 42°36'00"E), 2644 m. 24.06.2013, 2♂♂; Kayseri-Bağpınar (38°49'12"N, 35°37'56"E), 1097 m. 14.09.2006, 11♀, 7♂♂; -Gömeç (38°52'17"N, 35°39'43"E), 1094 m. 14.09.2006, 2♀♀, 1♂; -Pınarbaşı-Kızıldere (38°40'20"N, 36°15'27"E), 1457 m. 06.06.2007,

1♀;-Yemliha (38°52'02"N, 35°14'32"E), 992 m. 15.09.2006, 3♀♀, 2♂♂; -Yeşilhisar-Araplı (38°14'05"N, 35°03'13"E), 1349 m. 18.07.2007, 3♀♀, 2♂♂; Kırıkkale-Karakeçili (39°35'39"N, 33°22'46"E), 551 m. 11.07.2007, 1♀; -İnce-Paşayeri (41°40'40"E, 27°4'8"E), 150 m. 02.08.2000, 1♀; -Kızılıcadere (41°41'33"N, 27°18'48"E), 162 m. 02.06.1993, 1♀; Kırşehir-Kaman-Darıözü (39°30'21"N, 33°47'08"E), 902 m. 16.09.2006, 1♂; Nevşehir-Avanos-Saruhan (38°42'47"N, 34°54'47"E), 958 m. 06.06.2007, 2♀♀, 1♂; -Gülşehir-Gülyeri (38°44'55"N, 34°40'27"E), 910 m. 12.09.2006, 3♂♂; Niğde-Çiftlik (38°10'14"N, 34°29'17"E), 1562 m. 18.07.2007, 1♀; Sivas-Kızılıkavraz (39°46'20"N, 37°10'36"E), 1250 m. 27.05.2001, 1♀; -Merkez-Tepeönü (39°39'32"N, 37°08'01"E), 1306 m. 01.06.2007, 2♀♀; Tekirdağ-Çorlu (41°10'28"N, 27°48'20"E), 70 m. 12.08.1993, 1♀; Tokat-Almus-Gümeleönü (40°18'40"N, 37°7'10"E), 740 m. 02.09.2003, 2♀♀; Tunceli-Pülümür-Kangallı (39°27'10"N, 39°51'43"E), 1343 m. 27.08.2008, 1♀, 1♂.

Distribution: Palearctic, Oriental.

Hosts: Coleoptera, Gasterophilidae: *Magdalinus* sp. Diptera, Anthomyiidae: *Pegomya hyoscyami* (Panzer 1809). Tephritidae: *Euphranta connexa* (Fabricius, 1794). Lepidoptera, Crambidae: *Loxostege sticticalis* (Linnaeus, 1761). Elachistidae: *Elachista* sp.; Gelechiidae: *Scrobipalpa atriplicella* (Fischer von Röslerstamm, 1841); *Syncopacma coronillella* (Treitschke, 1833); *Teleiodes luculella* (Hübner, 1813). Gracillariidae: *Acrocercops brongniardella* (Fabricius, 1798); *Caloptilia alchimiella* (Stainton, 1850); *C. robustella* Jäckh, 1972; *C. rufipennella* (Hübner, 1796); *Gracillaria syringella* (Fabricius, 1794); *Parornix* sp.; *Phyllonorycter* sp.; *Phyllonorycter emberizaepennella* (Bouché, 1834); *P. ilicifoliella* (Duponchel, 1843); *P. nigrescentella* (Logan, 1851); *P. quercifoliella* (Zeller, 1839); *P. roboris* (Zeller, 1839). Lyonetiidae: *Leucoptera laburnella* (Stainton, 1851). Momphidae: *Mompha raschkiella* (Zeller, 1839); *Mompha terminella* (Humphreys & Westwood, 1845). Tineidae: *Agnathosia mendicella* (Denis & Schiffermüller, 1775). Tischeriidae: *Emmetia gaunacella* (Duponchel, 1843). Tortricidae: *Ancylis mitterbacheriana* (Denis & Schiffermüller, 1775); *Caloptilia betulicola* (M Hering, 1928); *Cydia cosmophorana* (Treitschke, 1835).

3.23. Genus *Rhyssalus* Haliday, 1833

Rhyssalus clavator Haliday, 1833

Material examined: Giresun-Duruoğlu (40°14'05"N, 38°7'00"E), 80 m. 05.07.2004, 1♀. 1♂.

Distribution: Western Palearctic.

3.24. Genus *Spathiomorpha* Tobias, 1976

Spathiomorpha varinervis Tobias, 1976

Material examined: Samsun-Salıpazarı-Derbentaltı (41°4'49"N, 36°49'35"E), 970 m. 03.07.2003, 2♂♂.

3.25. Genus *Spathius* Nees, 1819

**Spathius* (*Spathius*) *curvicaudis* Ratzeburg, 1844

Material examined: Adana-Osmaniye (36°59'59"N, 35°19'27"E), 26 m. 11.05.2000, 1♀; Denizli-Buharkent (37°57'42"N, 28°44'32"E), 173 m. 11.05.2000, 2♀♀; Kirkclareli-İğneada (41°53'33"N, 27°59'71"E), 10 m. 17.06.1987, 2♀♀; Sinop-Kabalı (41°52'42"N, 35°07'04"E), 70 m. 03.07.2001, 1♀.

Distribution: Western Palearctic.

Hosts: Coleoptera, Anobiidae: *Hedobia pubescens* (Olivier 1790); *Xestobium plumbeum* (Illiger, 1801). Buprestidae: *Agrilus biguttatus* (Fabricius, 1776); *A. pannonicus* (Piller & Mitterpacher, 1783); *A. roberti* (Chevrolat, 1837); *A. subauratus* (Gebler, 1833); *A. sulcicollis* (Lacordaire, 1835). Cerambycidae: *Callidium aeneum* (De Geer, 1775); *Obrium cantharinum* (Linnaeus, 1767); *Phymatodes pusillus* (Fabricius, 1787); *P. testaceus* (Linnaeus, 1758); *Pyrrhidium sanguineum* (Linnaeus, 1758). Curculionidae: *Magdalis carbonaria* (Linnaeus, 1758); *Scolytus intricatus* (Ratzeburg, 1837). Hymenoptera, Argidae: *Xiphydria camelus* (Linnaeus, 1758); *X. longicollis* (Geoffroy, 1785); *X. prolongata* (Geoffroy 1785); *X. pantherinus* (Savenius, 1825).

Spathius (*Spathius*) *erythrocephalus* Wesmael, 1838

Material examined: Samsun-Salıpazarı-Derbentli (41°4'49"N, 36°49'35"E), 970 m. 03.07.2003, 1♂.

Distribution: Western Palearctic.

Hosts: Coleoptera, Anobiidae: *Ochina ptinoides* (Marsham, 1802). Buprestidae: *Capnodis tenebrionis* (Linnaeus, 1758). Cerambycidae: *Callidiellum rufipenne* (Motschulsky, 1860); *Callidium aeneum* (De Geer, 1775); *Phymatodes testaceus* (Linnaeus, 1758). Curculionidae: *Crypturgus cinereus* (Herbst 1793); *Scolytus mali* (Bechstein & Scharfenberg, 1805). Lepidoptera, Tineidae: *Opogona* sp. Sesiidae: *Synanthedon vespermis* (Linnaeus, 1761). Hymenoptera, Xiphydriidae: *Xiphydria longicollis* (Geoffroy, 1785).

Spathius (*Spathius*) *exarator* (Linnaeus, 1758)

Material examined: Zonguldak-Devrek-Davulgâ (41°13'30"N, 31°57'34"E), 800 m. 29.06.2001, 2♂.

Distribution: Holarctic, Oceanic, Oriental.

Hosts: Coleoptera, Anobiidae: *Anobium inexpectatum* (Lohse, 1954); *A. punctatum* (DeGeer, 1774); *Dendrobiumpertinax* (Mulsant & Rey, 1863); *Dorcatoma* (*Pilosodorcatoma*) *setosella* (*setosella*) (Mulsant & Rey 1864); *D. dresdensis* (Herbst, 1792); *Ernobius mollis* (Linnaeus, 1758); *E. nigrinus* (Sturm, 1837); *Grynobius planus* (Fabricius, 1787); *Ochina ptinoides* (Marsham, 1802); *Ptilinus pectinicornis* (Linnaeus, 1758); *P. pectinicornis* (Linnaeus, 1758); *Ptinus* (*Ptinus*) *fur* (Linnaeus, 1758); *Scolytus fagi* Walsh 1867; *Stegobium paniceum* (Linnaeus 1761); *Xestobium rufovillosum* (De Geer, 1774); *X. rufovillosum* (De Geer, 1774). Buprestidae:

Agrilus ater (Linnaeus, 1767); *Anthaxia godeti* (Laporte de Castelnau & Gory, 1847); *A. quadripunctata* (Linnaeus, 1758); *Melanophila cyanea* (Fabricius, 1775); *M. guttulata* (Gebler, 1830). **Cerambycidae:** *Acanthocinus aedilis* (Linnaeus, 1758); *Callidium abdominalis* (Fabricius, 1787); *C. aeneum* (De Geer, 1775); *C. violaceum* (Linnaeus, 1758); *Clytus tropicus* (Panzer, 1795); *Exocentrus lusitanus* (Linnaeus, 1767); *Molorchus* (*Molorchus*) *minor* (Linnaeus, 1758); *Monochamus galloprovincialis* (Olivier, 1795); *Phymatodes pusillus* (*Pusillus*) (Fabricius, 1787); *P. testaceus* (Linnaeus, 1758); *Pogonocherus hispidus* (Linnaeus, 1758); *Rhagium inquisitor* (Linnaeus, 1758); *Semanotus undatus* (Linnaeus, 1758); *Stenostola ferrea* (Schrank, 1776); *S. ferrea* (Schrank, 1776); *Tetropium castaneum* (Linnaeus, 1758); *T. fuscum* (Fabricius, 1787); *T. gabrieli* (Weise, 1905); *T. gracilicorne* (Reitter, 1889). **Chrysomelidae:** *Chrysomela alni* (Linnaeus, 1758). **Curculionidae:** *Blastophagus piniperda* (Linnaeus, 1758); *Cryphalus tiliae* (sensu C.G. Thomson, 1865); *C. tibialis* (Lea, 1898); *Ernoporus fagi* (Fabricius, 1798); *Hylesinus fraxini* (Panzer, 1779); *Ips cembrae* (Heer, 1836); *I. sexdentatus* (Börner, 1767); *I. subelongatus* (Motchulsky, 1860); *I. typographus* (Linnaeus, 1758); *Magdalais* (*Magdalais*) *violacea* (Linnaeus, 1758); *Pentarthrum huttoni* (Wollaston, 1854); *Phloeosinus serrifer* (Wichmann, 1916); *Pissodes harcyniae* (Herbst, 1795); *P. nitidus* (Roelofs, 1874); *P. notatus* (Fabricius, 1787); *Pityogenes bidentatus* (Herbst, 1784); *P. bidentatus* (Herbst, 1784); *Pityophthorus micrographus* (Linnaeus, 1758); *Polygraphus subopacus* (Thomson, 1871); *Rhyncolus culinaris* (Hustache, 1931); *Rhynchaenus fagi* (Linnaeus, 1758); *R. pilosus* (Fabricius, 1781); *R. quercus* (Linnaeus, 1758); *R. salicis* (Linnaeus, 1758); *R. testaceus* (Mueller, 1776); *Scolytus multistriatus* (Marsham, 1802); *S. rugulosus* (Mueller, 1818); *S. scotylus* (Fabricius, 1775). **Tephritidae:** *Trupanea stellata* (Fuesslin, 1775). **Tortricidae:** *Cydia funebrana* (Treitschke, 1835); *C. strobilella* (Linnaeus, 1758). **Xiphydriidae:** *Xiphydria camelus* (Linnaeus, 1758); *X. prolongata* (Geoffroy 1785); *Xyela julii* (Brebisson 1818).

Spathius (*Spathius*) *polonicus* Niegabitowski, 1910

Material examined: Diyarbakır-Ergani-Çukurdere Köyü (38°5'42"N, 39°13'46"E), 1145 m. (Kuşburnu), 06.06.2007, 1♀; Diyarbakır-Ergani (38°16'09"N, 39°45'42"E), 872 m. 06.06.2007, 1♀.

Distribution: Western Palearctic.

Hosts: Coleoptera, Buprestidae: *Agrilus constantini* (Obenberger, 1927); *Agrilus viridis* (Linnaeus, 1758); *Anthaxia* sp.; *Coraebus florentinus* (Herbst, 1801); *Lampra mirifica* (Obenberger 1934); *Melanophila decastigma* (Fabricius, 1787); *M. Melanophila picta* (Pallas, 1773); *Sphenoptera davatchii* (Descarpentries, 1960); *Sphenoptera kaznakovi* (Jakovlev, 1899).

Spathius (*Spathius*) *radzayanus* Ratzeburg, 1848

Material examined: Adana-Osmaniye (36°59'59"N, 35°19'27"E), 26 m. 21.07.1980, 1♀; Adapazarı-Karasu

Harası (38°20'36"N, 38°03'35"E), 881 m. 08.07.1993, 1♀; Ankara-Elmadağ (39°55'15"N, 33°13'50"E), 1165 m. 16.09.2006, 1♀; Bursa-Cumalıķızlık (40°30'40"N, 29°18'38"E), 253 m. 18.09.1992, 1♀; Çanakkale-Küçükkyuyu (39°32'57"N, 26°36'34"E), 142 m. 04.11.2004, 1♀; Denizli-Buldan-Kadıköy (38°9'43"N, 28°45'32"E), 580 m. 19.09.1996, 1♀; Samsun-Salıpazarı-Astepe mevkii (41°4'49"N, 36°49'35"E), 600 m. 03.07.2003, 1♀.

Distribution: Palearctic, Oriental.

Hosts: Coleoptera, Buprestidae: *Agrilus auricollis* (Kiesenwetter, 1857); *A. biguttatus* (Fabricius, 1776); *A. viridis* (Linnaeus, 1758). **Anobiidae:** *Anobium abietis* (Fabricius 1792); *Ernobius nigrinus* (Sturm, 1837). **Cerambycidae:** *Exocentrus lusitanus* (Linnaeus, 1767); *Rhagium inquisitor* (Linnaeus, 1758). **Curculionidae:** *Hypothenemus obscurus* (Fabricius, 1801; syn. *Pissodes obscurus*); *Rhinocyllus conicus* (Frölich, 1792). **Scolitidae:** *Blastophagus piniperda* (Linnaeus, 1758). **Sesiidae:** *Synanthedon conopiformis* (Esper, 1782); *Synanthedon vespiformis* (Linnaeus, 1761). **Oligochaeta, Lumbricidae:** *Rhinocyllus depressicornis* (Smith, 1827).

**Spathius* (*Spathius*) *rubidus* (Rossi, 1794)

Material examined: Çanakkale-Küçükkyuyu (39°32'57"N, 26°36'34"E), 14 m. 26.10.2010, 1♀; Erzurum-Pasinler-Övenler (39°59'16"N, 41°34'38"E), 1740 m. 17.08.2013, 1♀; Hatay-Dörttyol (36°50'43"N, 36°13'22"E), 71 m. 30.07.1983, 1♀; İstanbul-Sarıyer Bilezik Çiçiftliği (41°10'53"N, 29°21'18"E), 35 m. 14.06.2001, 1♀; İzmir-Aliağa (38°47'41"N, 26°58'13"E), 10 m. 02.06.2002, 1♀; Kastamonu-Tosya-Ekinci (41°1'6.64"N, 34°22'14"E), 1000 m. 31.08.2002, 1♀; Kırklareli-Dereköy-Kösayaşı (41°56'43"N, 27°21'57"E), 500 m. 29.07.1987, 1♀; Tokat-Almus-Gömelönü (40°18'40"N, 37°7'10"E), 740 m. 02.09.2003, 1♂; Trabzon-Maçka-Dikkaya (40°42'15"N, 39°29'53"E), 1016 m. 05.08.2005, 1♂; Tokat-Niksar (40°35'12"N, 36°57'72"E), 339 m. 05.07.2003, 1♀.

Distribution: Holarctic.

Hosts: Coleoptera, Anobiidae: *Anobium punctatum* (DeGeer, 1774); *A. (Hadrobregmus) rufipes* Fabricius, 1792; *Ernobius longicornis* Sturm, 1837; *Hedobia imperialis* (Linnaeus, 1767); *Ochina ptinoides* (Marsham, 1802). **Bostrichidae:** *Scobicia chevrieri* (Villa & Villa, 1835); *Sinoxylon exdentatum* (Olivier, 1790); *Xylopertha praeusta* (Germar, 1817); *X. retusa* (Olivier, 1790). **Buprestidae:** *Agrilus angustulus* (Illiger, 1803); *A. auricollis* Kiesenwetter, 1857; *A. laticornis* (Illiger, 1803); *A. viridis* (Linnaeus, 1758); *A. biguttatus* (Fabricius, 1776); *A. convexicollis* Redtenbacher 1849; *Melanophila picta* (Pallas, 1773). **Cerambycidae:** *Leiopus nebulosus* (Linnaeus, 1758); *Phymatodes testaceus* (Linnaeus, 1758); *Pogonocherus fasciculatus* (DeGeer, 1775); *Pyrrhidium sanguineum* (Linnaeus, 1758); *Saperda populnea* (Linnaeus, 1758); *Tetrops praeustus* (Linnaeus 1758); *T. starki* Chevrolat,

1859. Chrysomelidae: *Agelastica alni* (Linnaeus, 1758). Colydiidae: *Synchita mediolanensis* Villa & Villa 1833. Curculionidae: *Blastophagus minor* (Hartig, 1834); *Caphoborus minimus* (Fabricius, 1798); *Ceutorhynchus quadridens* Panzer, 1795; *Conophthorus coniperda* (Schwarz, 1895); *Cryphalops tiliae* (Panzer, 1793); *Crypturgus cinereus* (Herbst, 1793); *Hylesinus fraxini* (Panzer 1779); *Ips acuminatus* (Gyllenhal, 1827); *I. latidens* (LeConte, 1874); *I. lecontei* Swaine, 1924; *I. subelongatus* Motchulsky, 1860; *I. lecontei* Swaine, 1924; *Lixus (Eulixus) bidens* Capiomont, 1874; *Magdalalis carbonaria* Stephens, 1831; *M. cerasi* (Linnaeus, 1758); *Magdalalis (Magdalalis) frontalis* (Gyllenhal, 1827); *M. (M.s.) rufa* Germar, 1824; *M. ruficornis* (Linnaeus, 1758); *Phloeosinus bicolor* (Brulle, 1832); *P. thujae* (Perris, 1855); *P. scarabaeoides* (Bernard, 1788); *Pissodes castaneus* (De Geer, 1775); *P. Pissodes notatus* Fabricius, 1787; *P. Pissodes validirostris* (Sahlberg, 1834); *Pityogenes bidentatus* (Herbst, 1784); *P. calcaratus* (Eichhoff, 1878); *P. chalcographus* (Linnaeus, 1761); *P. quadridens* (Hartig, 1834); *Pityophthorus puberulus* (LeConte, 1868); *Rynchaenus (R.) quercus* (Linnaeus, 1758); *Scolytus intricatus* (Ratzeburg, 1837); *S. multistriatus* (Marsham, 1802); *S. opacus* Blackman, 1934; *S. pygmaeus* (Fabricius, 1787); *S. rugulosus* (Müller, 1818); *S. scotylus* (Fabricius 1775). Hymenoptera, Xiphydriidae: *Xiphydria longicollis* (Geoffroy, 1785); *X. prolongata* (Geoffroy, 1785).

4. Discussion

The present study on the Doryctinae fauna of Turkey revealed the existence of 58 species belonging to 25 genera, of which 52 species and 20 genera are recorded for the first time from Turkey. The present study is the first summary of all recorded species of the subfamily Doryctinae in Turkey. Previous studies were conducted in forest parts of Turkey and 7 species were recorded.

The most common species is *Hormius moniliatus* (Nees) and it was collected from 125 localities. *H. moniliatus* (Nees) has been recorded from Holarctic, Nearctic, Oceanic, and Oriental regions and is known to have 15 host species (forest pests). The second common species, *Rhysipolis meditator* (Haliday), which was

collected from 44 localities, is distributed in Palearctic and Oriental regions. Each of 21 species was recorded from 1 locality. They can therefore be considered as rare species.

Doryctine species are parasitoids of bark beetles, which are very important forest pests. They are important population regulatory factors of species of Coleoptera, Diptera, Lepidoptera, Hymenoptera, and Diptera (Shaw and Huddleston, 1991).

The following doryctine species are known to have more host species: the number of host species of *Spathius* (S.) *exarator* (Linnaeus) is 72, of *S. (S.) rubidus* (Rossi) 62, and of *Ecpylus* (E.) *silesiacus* (Ratzeburg, 1848) is 63. One of the 60 host species of *Doryctes* (*D.*) *leucogaster* (Nees) is *Opilo domesticus* (Sturm) (Aranaea, Theridiidae). *Rhinocyllus depressicornis* (Smith) (Oligochaeta, Lumbricidae) is a host of *Spathius* (*Spathius*) *polonicus* Niegabitowski.

The zoogeographical distributions of species of Doryctinae are as follows: 45 species are recorded only from Palearctic (5 species from the Eastern Palearctic, 17 species from the Western Palearctic); 8 species from Palearctic and Oriental; 2 species from Holarctic, Oceanic, and Oriental; 1 species from Holarctic and Oriental; and 1 species from Holarctic, Neotropic, and Oriental regions.

The distribution area of *Colastes* (*Colastes*) *braconius* Haliday is the Palearctic region. It was introduced into many parts of America to be used in biological control of forest pests (USA-Delaware; USA-New Jersey; USA-Pennsylvania) (Hendrickson and Barth, 1979).

Coeloides forsteri Haeselbarth was obtained for the first time from an *Ips sexdentatus* larvae host.

Acknowledgments

This research was supported by the Scientific and Technological Research Council of Turkey (TÜBİTAK, Projects 192T057, 196T029, 103T176, 106T588, 111T416) and the Trakya University Scientific Research Project Unit (TÜBAP Projects 462, 553, 740). The author thanks both institutions for their financial support, as well as Dr Belokobylskij from the Zoological Institute of the Russian Academy of Sciences (St Petersburg) for the confirmation of the species belonging to Doryctinae.

References

- Barbalho SM, Penteado-Dias AM, Marsh PM (1999). Description of new genera from Brazil in the tribes Heterospilini and Spathiini with similar wing venation (Hymenoptera: Braconidae, Doryctinae). *J Hymenopt Res* 8: 139–153.
- Belokobylskij SA (1986). [A review of the Palaeartic species of the genera *Pambolus* Hal. and *Dimerus* Ruthe (Hymenoptera, Braconidae)]. *Trudy Zoologicheskogo Instituta Leningrad* 159: 18–37 (in Russian).
- Belokobylskij SA (1990). Review of braconid wasps of the genus *Rhaconotus* Ruthe (Hymenoptera, Braconidae) of the Palearctic Region. *Entomol Obozr* 69: 144–163.
- Belokobylskij SA (1992). On the classification and phylogeny of the Braconid wasps subfamilies Doryctinae and Exothecinae (Hymenoptera, Braconidae). I. Classification. *Entomol Rev* 72: 109–137.

- Belokobylskij SA (1993). On the classification and phylogeny of Braconidae wasps of subfamilies Doryctinae and Exothecinae (Hymenoptera, Braconidae). Part II. Phylogeny. Entomol Rev 73: 1–27.
- Belokobylskij SA (1994a). A new tribe of the subfamily Doryctinae from Papua New Guinea (Hymenoptera: Braconidae). Zoosyst Rossica 3: 141–145.
- Belokobylskij SA (1994b). Obzornaezdnikov-brakonidpodsemejstva Doryctinae, Exothecinae (Hymenoptera, Braconidae) Dalnego Vostoka, Vostotsnoi Sibiri Sopredelnychterritoriij. Trudy Zapov Dahursky, Kiev [= Hymenoptera Insects of Siberia and Far East]. Memoirs of the Dahursky Nature Reserve 3: 5–77.
- Belokobylskij SA (1995). Two new genera and two new subgenera of the subfamilies Exothecinae and Doryctinae from the Old World (Hymenoptera: Braconidae). Zool Meded Leiden 69: 37–52.
- Belokobylskij SA (1998). On braconid wasps of the genus *Ontsira* Cameron (Hymenoptera, Braconidae) of East Asia. Entomol Rev 78: 556–564.
- Belokobylskij SA (2001). [New taxa of the braconid subfamilies Doryctinae and Exothecinae (Hymenoptera, Braconidae) from west Palearctic.]. Entomologicheskoe Obozrenie 80: 451–471 (in Russian).
- Belokobylskij SA (2003). The species of the genus *Spathius* Nees, 1818 (Hymenoptera: Braconidae: Doryctinae) not included in the monograph by Nixon (1943). Ann Zool 53: 347–488.
- Belokobylskij SA (2004a). New genus and new subgenus of subfamily Doryctinae (Hymenoptera: Braconidae) from the old world fauna. Ann Soc Entomol Fr 40: 199–204.
- Belokobylskij SA (2004b). Two new taxa of subtribe Rhaconotina (Hymenoptera: Braconidae, Doryctinae, Doryctini) from Africa, with a key to subtribe genera. Ann Soc Entomol Fr 40: 205–210.
- Belokobylskij SA, Chen X (2004). The species of the genus *Rhaconotus* Ruthe, 1854 (Hymenoptera: Braconidae: Doryctinae) from China with a key to species. Ann Zool 54: 319–359.
- Belokobylskij SA, Chen X (2006). *Hecabolomorpha* n. gen., a new Asian genus from the tribe Hecabolini (Hymenoptera: Braconidae: Doryctinae). Ann Soc Entomol Fr 42: 107–111.
- Belokobylskij SA, Iqbal M, Austin AD (2004a). Systematics, distribution and diversity of the Australian doryctine wasps (Hymenoptera, Braconidae, Doryctinae). Records of the South Australian Museum Monograph Series 8: 1–150.
- Belokobylskij SA, Maeto K (2006). A new species of the genus *Parachremylus* Granger (Hymenoptera: Braconidae), a parasitoid of *Conopomorpha* lychee pests (Lepidoptera: Gracillariidae) in Thailand. J Hymenopt Res 15: 181–186.
- Belokobylskij SA, Maeto K (2008). Doryctinae (Hymenoptera: Braconidae) of Ogasawara Islands (Japan). Ann Zool 58: 125–166.
- Belokobylskij SA, Maeto K (2009). Doryctinae (Hymenoptera, Braconidae) of Japan. (Fauna Mundi.Vol 1) Warsaw, Poland: Warshawskia Drukarnia Naukova.
- Belokobylskij SA, Quicke DLJ (2000). Seven new genera of the subfamily Doryctinae (Hymenoptera: Braconidae) from the Old World. J Hymenopt Res 9: 111–141.
- Belokobylskij SA, Samartsev KG (2011). First record of the tribe Holcobraconini and the genus *Zombrus* Marshall, 1897 (Hymenoptera: Braconidae: Doryctinae) in Europe. Zoosyst Rossica 20: 310–318.
- Belokobylskij SA, Tang P, He J, Chen X (2012). The genus *Doryctes* Haliday, 1836 (Hymenoptera: Braconidae, Doryctinae) in China. Zootaxa 3226: 46–60.
- Belokobylskij SA, Zaldivar-Riverón A, Quicke DLJ (2004b). Phylogeny of the genera of the parasitic wasps subfamily Doryctinae (Hymenoptera: Braconidae) based on morphological evidence. Zool J Linn Soc 142: 369–404.
- Belokobylskij SA, Žikić V (2009). New data on cyclostome braconid subfamilies Doryctinae, Exothecinae, Rogadinae and Braconinae (Hymenoptera: Braconidae) of Serbia and neighbouring territories. Acta Entomol Serb 14: 65–71.
- Beyarslan A, Aydogdu M (2013). Additions to the rare species of Braconidae fauna (Hymenoptera: Braconidae) from Turkey. Mun Ent Zool 8: 369–374.
- Fischer M (1970). Eineneue Spathius-Art ausdem Iran (Hym. Braconidae). Ann Soc Entomol Fr 6: 705–708 (in German).
- Fischer M (1980). Taxonomische Untersuchungenüber Doryctinae aus der Odontobracon- Verwandtschaft (Hymenoptera, Braconidae). Ann Naturhist Mus Wien 83: 547–572 (in German).
- Ghahari H, Gadallah NS, Cetin Erdogan O, Hedqvist KJ, Fischer M, Beyarslan A, Ostovan H (2009). Faunistic note on the Braconidae (Hymenoptera: Ichneumonoidea) in Iranian cotton fields and surrounding grasslands. Egypt J Genet Cytol Bio Pest Cont 19: 115–118.
- Gomes SAG, Penteado-Dias AM (2007). Description of a new genus of Doryctinae wasps (Hymenoptera: Braconidae) from Brazil. Zool Meded (Leiden) 81: 115–119.
- Hedqvist KJ (1976). New species of *Spathius* Nees, 1818 and a key to the species of Europe and Canary Islands (Hym. Ichneumonoidea, Braconidae). Eos 51: 51–63.
- Hendrickson RM, Barth SE (1979). Introduced parasites of *Agromyza frontella* (Rondani) in the USA. J N Y Entomol Soc 87: 167–174.
- Kula R (2009). Review of the new world species of *Coiba* Marsh (Hymenoptera: Braconidae: Doryctinae), including descriptions of two new species, new distribution records, and a key to species. Proc Entomol Soc Wash 111: 183–198.
- Loni A, Hart RS, Lucchi A (2012). First record of *Zombrus bicolor* (Enderlein) (Hymenoptera, Braconidae, Doryctinae) in Western Europe. Zookeys 219: 89–91.
- Mancini D, Priore R, Battaglia D, van Achterberg C (2003). *Caenopachys hartigii* (Ratzeburg) (Hymenoptera: Braconidae: Doryctinae) confirmed for Italy, with notes on the status of the genus *Caenopachys* Foerster. Zool Meded (Leiden) 77: 459–470.

- Marsh PM (1993). Descriptions of new Western Hemisphere genera of the subfamily Doryctinae (Hymenoptera: Braconidae). Contributions of the American Entomological Institute 28: 1–58.
- Marsh PM (1997). Subfamily Doryctinae. In: Wharton RA, Marsh PM, Sharkey MJ, editors. Manual of the New World Genera of the Family Braconidae (Hymenoptera): Subfamily Doryctinae. Washington, DC, USA: Special Publications of the International Society of Hymenopterists, pp. 206–233.
- Marsh PM (2002). The Doryctinae of Costa Rica (excluding the genus *Heterospilus*). Mem Am Entomol Inst 70: 1–229.
- Marsh PM, Melo GAR (1999). Biology and systematics of new world *Heterospilus* (Hymenoptera: Braconidae) attacking Pemphredoninae (Hymenoptera: Sphecidae). J Hymenopt Res 8: 13–22.
- Martinez JJ, Zaldivar-Riverón A (2013). Seven new species of *Allorhogas* (Hymenoptera: Braconidae: Doryctinae) from Mexico. Revista Mexicana de Biodiversidad 84: 117–139.
- Matthews RW (1970). A revision of the genus *Spathius* in America north of Mexico (Hymenoptera: Braconidae). Entomol Soc Am 4: 1–86.
- Quicke DLJ (1987). The Old World genera of braconine wasps (Hymenoptera: Braconidae). J Nat Hist 21: 43–157.
- Quicke DLJ (1997). Parasitic Wasps. London, UK: Chapman & Hall.
- Quicke DLJ, Ficken LC, Fitton M (1992). New diagnostic ovipositor characters for doryctine wasps (Hymenoptera, Braconidae). J Nat Hist 26: 1035–1046.
- Ramirez BW, Marsh PM (1996). A review of the genus *Psenobolus* (Hymenoptera: Braconidae) from Costa Rica, an inquiline fig wasp with brachypterous males, with descriptions of two new species. J Hym Res 5: 64–72.
- Rastegar J, Sakenin H, Khodaparast S, Havaskary M (2012). On a collection of Braconidae (Hymenoptera) from East Azerbaijan and vicinity, Iran. Calodema 226: 1–4.
- Shaw MR, Huddleston T (1991) Classification and Biology of Braconid Wasps. Handbooks for the Identification of British Insects, Vol. 7. St Albans, UK: Royal Entomological Society.
- Shaw SR, Edgerly JS (1985). A new braconid genus (Hymenoptera) parasitizing webspinners (Embiidina) in Trinidad. Psyche 92: 505–511.
- Schimitschek E (1939). Die Lebensweise des ryocoetes minor. Forstarchiv 15: 271–274 (in German).
- Schimitschek E (1941). Forstinsekten der Tuerkei und ihre Umwelt. Prague: Volk und Reich (in German).
- Schimitschek E (1944). Forstinsekten der Tuerkei und ihre Umwelt. Prague: Volk und Reich (in German).
- Shenefelt RD, Marsh PM (1976). Braconidae. Doryctinae. In: Ferrière C, van der Vecht J, editors. Hymenopterorum Catalogus (nova editio) Pars 13. The Hague, Netherlands: Uitgeverij Dr. W. Junk, pp. 1263–1424.
- Tang P, Belokobylskij SA, He J, Chen X (2013). *Heterospilus* Haliday, 1836 (Hymenoptera: Braconidae, Doryctinae) from China with a key to species. Zootaxa 3683: 201–246.
- Telenga NA (1941). Family Braconidae, subfamily Braconinae (continuation) and Sigalphinae. Fauna USSR. Hymenoptera 5(3). Moscow: Akademiya Nauk SSSR.
- Tobias VI (1986). Subfamily Doryctinae. In: Medvedev GS, editor. Keys to the Insects of the European Part of the USSR: Subfamily Doryctinae. Volume III, Hymenoptera Part IV. New Delhi: Amerind Publishing Co. Pvt. Ltd., pp. 28–117.
- van Achterberg C (1993). Illustrated key to the subfamilies of the Braconidae (Hymenoptera: Ichneumonoidea). Zool Meded (Leiden) 283: 1–189.
- van Achterberg C (2003). The West Palaearctic species of the genera *Gildoria* Hedqvist and *Platyspathius* Viereck, with keys to the species (Hymenoptera: Braconidae: Doryctinae). Zool Meded (Leiden) 77: 267–290.
- Wharton RA, Hanson PE (2005). Biology and evolution of braconid gall wasps,. In: Ramon A, Schaefer CW, Withers TM, editors. Biology, Ecology and Evolution of Gall-Inducing Arthropods. Enfield, NH, USA: Science Publishers, pp. 495–505.
- Yu DS, van Achterberg C, Horstmann K (2012). Ichneumonoidea 2011 (Biological and taxonomical information). Taxapad Interactive Catalogue. Available at www.taxapad.com.
- Zaldivar-Riverón A, Belokobylskij SA (2009). The parasitic wasp genus *Hecabolus* Curtis (Hymenoptera: Braconidae: Doryctinae), with the description of a new species from Mexico. Revista Mexicana de Biodiversidad 80: 419–429.
- Zaldivar-Riverón A, Belokobylskij SA, Leon RV, Briceno GR, Quicke DLJ (2008). Molecular phylogeny and historical biogeography of the cosmopolitan parasitic wasp subfamily Doryctinae (Hymenoptera: Braconidae). Invertebr Syst 22: 345–363.